

ENVIRONMENT

Title: Swine Manure Applications for Soybean Production – Environmental and Pathological Implications – **NPB #08-052**

Investigator: Robert W. Mullen, Brian McSpadden-Gardner, Keith Diedrick, Rosa Raudales, Jon Rausch, and Clay Dygert

Institution: Ohio State University/Ohio Agricultural Research and Development Center

Date Submitted: February 25, 2010

INDUSTRY SUMMARY

The application of manure to soybeans has been identified as a potential environmental threat due to the fact that soybeans are legumes and do not require nitrogen supplementation. Thus, the application of animal manures may pose a risk to water quality. Our goal in this research project was to ascertain the actual risk of nitrogen loss by measuring nitrate-N accumulation in the soil and plant tissues throughout the growing season after a swine manure application (and corresponding commercial fertilizer treatments) and determine whether or not these applications present an environmental concern. Additionally, we were attempting to determine some potential guidelines in recommending a safe application rate of manure N. Our results indicate that application of manure N or commercial N at reasonable agronomic rates (<120 pounds of N per acre) should be of little environmental concern. This is supported by a lack of nitrate-N accumulation in the top 12” of soil, and a typical increase in tissue N accumulation when supplemental N is provided. Soybeans will accumulate soil N when it is supplied by manure or synthetic commercial sources. Higher rates of N (>180 pounds N per acre) from either source may pose an environmental risk in terms of excess soil nitrate-N and therefore should be discouraged.

ABSTRACT

Recent environmental pressure has been placed on the practice of applying animal manures to fields going into soybean production because of the perceived over-application of nitrogen to a legume crop that does not require nitrogen fertilization. The objective of this study was to evaluate the impact of manure application to soybean fields on soil nitrate accumulation and soybean crop productivity. Additionally, this work was conducted to determine the impact of manure application on soilborne pathogens specifically soybean cyst nematode. A single field experiment was established at the Western Research Station near South Charleston, OH in the spring of 2007. Liquid swine manure was applied at three different nitrogen (N) rates (based upon manure analysis and estimated availability) using two application methods (surface application or injection). The surface application treatments were either incorporated by tillage or left on the soil surface. Commercial fertilizer treatments (same N rates) were also included as positive controls. After soybean planting, soil samples were collected to measure soil nitrate to a depth of 2 feet at three different times during the growing season and after crop harvest. At two different growth stages (V4 and R1) soybean roots were collected and analyzed to

These research results were submitted in fulfillment of checkoff-funded research projects. This report is published directly as submitted by the project's principal investigator. This report has not been peer-reviewed.

For more information contact:

National Pork Board • PO Box 9114 • Des Moines, IA 50306 USA • 800-456-7675 • Fax: 515-223-2646 • pork.org

determine soybean cyst nematode infection levels. Even though N application did result in higher soil nitrate levels for some treatments, applications rates that were lower (less than or equal to 120 lb N/acre) represent a lower risk of possible nitrate loss. This is especially true later in the growing season. Early in the growing season application of N supplying fertilizers did result in higher nitrate levels. This is expected when plant growth is progressing slowly and nutrient demand from the soil is low. Nitrogen supplied via manure and commercial fertilizer resulted in larger N uptake than the controls at the later stages of growth. This reveals that despite the fact that soybeans can fix their own N, soybeans will absorb soil inorganic N if it is available. This work also reveals that fertilization may decrease soybean cyst nematode infection although at this point the mechanism is not fully understood.

INTRODUCTION

Soybeans are leguminous plant species capable of fixing nitrogen (N) from the atmosphere to support their growth and development. Despite the fact that soybeans can fix their own N, in a soil environment rich in inorganic N they will scavenge for available N decreasing the level of root nodulation (Schmidt et al., 2000; Hesterman and Isleib, 1991). Studies conducted to measure the impact of N fertilization of soybeans reveals that crop response to commercial N fertilizer does not result in increased crop production unless soil conditions inhibit rhizobium infection (low soil pH, cool, wet soils, etc.), which are the bacteria responsible for N fixation (Reese Jr., and Buss, 1992; Schmidt et al., 2000; Osborne and Riedell, 2006). Because soybeans are unlikely to benefit from N fertilization, most land grant universities do not recommend application of N-containing fertilizers to soybeans because it is not economical to do so (unless those extraneous soil conditions exist). Manure application for soybean production can provide needed phosphorus, potassium, and additional nutrients, but because of the environmental risks associated with excessive application of N, state and federal agencies have considered encouraging producers not to apply manure on soybean production ground. This could place more pressure on swine operations to identify available land for recycling manure nutrients.

Previous work has shown that soybeans can take up soil inorganic nitrogen from both manure and commercial forms of nitrogen. The most extensive work was conducted in Minnesota, but this experiment relied on post-harvest soil profile nitrate analysis to determine residual nitrate to a depth of four feet (Schmidt et al., 2000). Ohio production fields are extensively tiled (drained) to an average depth of 30 to 36 inches, and thus a single post-harvest estimate of profile nitrate may be misleading. This is especially true if the growing season has been exceptionally wet.

Organic soil amendments can have positive impacts on soilborne pathogens and plant health (Hoitink and Boehm 1999, Lazarovits 2001). In soybeans, several researchers have conducted investigations into the effects of manure applications on crop health and found that results are dependent upon the timing and amount of organic amendment applied. For example, application of compost 3 to 6 months prior to soybean planting resulted in decreased *Phytophthora* damage and increased yield in moderately susceptible cultivars (Hoitink and Schmitthenner, 1988). Swine manure has also shown some suppressive ability on soybean cyst nematode egg hatch attributed to high salt content and high concentration of organic chemicals, though SCN egg counts were increased in the fall due to manure application (Reynolds et al., 1999). However, high rates of swine manure application have also been shown to aggravate other soilborne disease problems such as white mold (Schmidt et al 2001). Thus, it is important to systematically determine how timing and rate affect disease pressure so as to not negatively offset the nutrient benefits of such applications.

OBJECTIVES

The objective of this study was to evaluate the impact of manure application to soybean fields on soil nitrate accumulation and soybean crop productivity. Additionally, this work was conducted to determine the impact of manure application on soilborne pathogens specifically soybean cyst nematode.

MATERIALS AND METHODS

A single field experiment was established in the spring of 2007 at the OARDC Western Research Station near South Charleston, OH. Swine manure was applied at three N rates 60, 120, and 180 lb/acre assuming that one third of the organic fraction was considered plant available and the ammonium-N manure

fraction was considered 75% plant available. Manure samples were collected from the pit during and after application and used to represent the average manure analysis (Table 2). Historical manure analysis was used to determine manure application rates for 2007 (Table 1), and 2007 manure analysis was used for the 2008 applications (Table 2). Liquid swine manure was applied either as a surface treatment or subsurface injected using Dietrich shanks to a depth of 4-5 inches. All plots that received subsurface injected liquid swine manure were tilled, and the surface application treatments were either no-tilled or tilled. Tillage comprised of a single tandem offset disk pass and a single pass of a finishing tool. In 2008, since the previous year's injected manure required tillage, we abandoned that plot and added another suite of treatments where commercial fertilizer could be surface applied without tillage. Liquid swine manure was applied to the soil surface treatments on April 10, 2007 and April 29, 2008. Due to rainfall on April 11, 2007 the subsurface treatments were applied on April 18, 2007 and tillage was done on April 20, 2007. A randomized complete block experimental design was used with four replications. Each plot was 10 ft by 30 ft. Phosphorus and potassium were supplied at rates equivalent to the amount of P and K supplied by the manure application. Urea, triple-superphosphate, and potash were the products used for N, P, and K, respectively, as the synthetic commercial treatments.

Soil samples were collected at four dates in 2007 – June 1, June 28, July 18, and September 25, and at three dates in 2008 – July 3, August 7, and October 30. Tissue samples were collected in 2007 on June 28 and July 18 (that corresponded with soybean vegetative stage V4 and R1, respectively). Tissue samples were collected in 2008 on July 3 and August 7 (that corresponded with soybean vegetative stage V4 and R1, respectively). Soil samples were collected to a depth of 24" if sampling conditions allowed and to a depth of 12" otherwise. Soil ammonium-N analysis was done using the steam distillation method, and nitrate-N analysis was done using an ion selective electrode (ISE). Tissue N concentration was determined using dry combustion. To ascertain the impact of soybean cyst nematode infection of plant roots, root samples were collected when soybean tissue was collected. Root samples were partitioned and tap root weight and total root weight were determined. Roots were visually inspected to determine SCN infection levels. Statistical analysis to determine differences between treatments was done using single degree of freedom contrasts in PROC GLM in SAS (SAS, 1998).

RESULTS AND DISCUSSION

Soybean Biomass

In 2007, soybean biomass measured at V4 was only increased above the corresponding control when swine manure was injected below the soil surface (Table 4). That application method also resulted in a linear increase in soybean biomass with increasing N application rate. In 2008, soybean biomass measured at V4 was only increased above the control when commercial urea was surface applied and incorporated (Table 13). Interestingly, that application method did not result in a linear increase in soybean biomass. This suggests that incorporation of the urea provided adequate N unlike the surface application without incorporation, but the higher N rates did not result in increased biomass. The absence of a significant response to applied swine manure indicates that not all the predicted mineralization of N occurred. It is also likely an artifact of the variation in N supply based on manure analysis.

In 2007, soybean biomass measured at R1 was increased above the corresponding control when manure or commercial fertilizer was surface applied and subsequently incorporated via tillage (Table 5). The only application method that did not result in a linear increase in soybean biomass was the swine manure surface application with tillage. These results may be explained by the positional availability of the applied nutrients due to tillage. Application methodologies that placed the nutrient closer to the root zone resulted in improved biomass accumulation. This response was unlikely due to the actual N application, but more likely attributable to the P and K added since soil test levels were at or below established critical levels (Table 3). In 2008, soybean biomass was only significantly different than the control when manure was surface applied and incorporated and when commercial urea was applied and incorporated (Table 14). This effect is not solely attributable to tillage since there was also a significant linear increase in biomass as N application rate increased.

Soybean Tissue Nitrogen Concentration

In 2007, soybean tissue N concentration measured at V4 increased linearly as N application rate increased for both the injected manure treatments and the commercial fertilizer treatments (Table 4). The lack of a significant difference between the control and the treatments means and a corresponding linear increase in tissue N concentration reveals that only at the high application rates were differences noted. In 2008, tissue N concentration was only increased above the control when with manure or commercial urea was surface applied and incorporated (Table 13). This is likely an indication that the surface applied manure and urea that was not incorporated was lost via volatilization.

In 2007, soybean tissue N concentration measured at R1 was increased above the corresponding control when manure was injected and when manure was surface applied and no tillage was done (Table 5). Subsequently, those same two application methods resulted in a linear increase in soybean tissue nitrogen concentration. This observation is slightly different that what was seen at the V4 stage measurements when the commercial fertilization treatments resulted in an increase in tissue N concentration. In 2008, soybean tissue concentration was increased above the control when manure was surface applied either with or without tillage (Table 14). Linear increases were observed when both manure and commercial urea were surface applied and incorporated.

Soybean Nitrogen Uptake

In 2007, soybean N uptake measured at V4 was increased above the control when manure was injected below the soil surface (Table 4). A linear increase in N uptake was noted when manure was injected or commercial fertilizer was surface applied and incorporated. Again, the lack of increased N uptake above the control and subsequent linear increase with increase N rate indicates that N uptake was only affected at the higher N application rates. The lack of response at this earlier growth stage is not surprising considering the small amount of biomass and nutrient accumulated at such early growth stages. In 2008, soybean N uptake was only different from the control when commercial urea was applied and incorporated (Table 13). This is an indication that we are underestimating N availability from the manure, or that mineralization of the organic-N fraction was occurring slower than estimated.

In 2007, soybean N uptake measured at R1 was increased above the control for all application methods except when manure was surface applied and incorporated with tillage (Table 5). This may be due to positional availability of nutrients applied or the lack of when tillage was not done. All application methods resulted in a linear increase in soybean N uptake with increasing N application rate. This indicates that application of inorganic-N (or organic-N that can readily be converted to inorganic-N) results in greater N uptake by the soybean crop, reinforcing that application of inorganic-N to a leguminous crop species will be readily be taken up in lieu of N fixation. In 2008, soybean N uptake was increased above the control when manure was surface applied (regardless of tillage) and when commercial urea was surface applied and incorporated (Table 14). The lack of significant N uptake when commercial urea was applied likely is an indication of volatilization losses of ammonia-N.

Soybean Yield

In 2007, soybean yield was increased above the control for all treatments (Table 6). A linear increase in yield with increasing N application rate was only noted for the injected manure treatment. In all likelihood, the N application itself was not the factor that resulted in increased yield. A more plausible explanation is that the manure and commercial treatments provided P and K to a soil that was near current established critical levels (especially K). This is supported by visual assessments of treatments on July 18 when it was noted that control treatments were showing visual symptoms of K deficiency (quite severe in some plots). Dry conditions during the summer months contributed to this condition as well, but supplementation either with swine manure or commercial fertilizer decreased or removed any signs of K stress. In 2008, soybean yield was improved above the control when manure was supplied (regardless of tillage) and when commercial fertilizers were supplied and incorporated (Table 18). As was noted in 2007, the likely reason for a yield increase is most likely attributable to supplementation of P and K due to low levels of both. The lack of yield improvement when manure was

injected on 30" centers and when commercial fertilizer was surface applied may speak to poor positional availability.

Soybean Cyst Nematode Root Counts

Tap root weight and total root weight were only increased above the control in a linear fashion with the application of commercial fertilizer for the first sampling date (Table 7). Nematodes infection was decreased below the control in a linear fashion when manure was surface applied and incorporated with tillage (Table 7). At the second sampling period, tap root weight and total root weight were not affected by treatment. Nematode infection level was decreased below the control treatments when manure was surface applied and incorporated with tillage or when commercial fertilizer was supplied (Table 8).

Soil Nitrate

In 2007, soil nitrate-N levels increased linearly at the 0-12" depth as N application rate increased across all application methodologies for the first soil sampling time (Table 9). With the exception of the injected manure treatment, all applications of nitrogen increased soil nitrate-N levels above that of the control. Soil nitrate-N levels were increased above the control for the 12-24" sampling depth when manure was surface applied and not incorporated with tillage and when commercial fertilizer was surface applied (Table 9). Those two treatments also showed a linear trend with increasing application rate. In 2008, soil nitrate levels increased linearly across most application methods and N sources at the 0-12" depth with the exception being the commercial urea surface applied and not incorporated (Table 15). Even though nitrate-N levels increased in a linear fashion does not necessarily indicate greater risk of accumulation of excess nitrate-N compared to the controls. Comparison of the manure injected and manure surface applied without tillage with the control reveals statistically similar nitrate-N levels. Nitrate-N in the 12-24" depth were only increased above the control when manure and commercial urea were surface applied and incorporated (Table 15). As was observed in 2007, analysis of the first sampling date shows little nitrate-N accumulation at the 0-12" depth if N application rates are close to soybean N removal rates. This was not true when commercial urea was applied and incorporated. Even less nitrate-N accumulates at the 12-24" level (as would be expected) except at the higher N rates when manure or commercial fertilizer is surface applied and incorporated.

At the second sampling date in 2007, soil nitrate levels in the 0-12" were increased above the control when manure was injected and when commercial fertilizer was surface applied and incorporated with tillage (Table 10). A linear increase in nitrate levels was observed with increasing N application rate for all treatments except when manure was applied to the surface and incorporated. At the 12-24" depth a linear increase in soil nitrate was noted when manure was injected below the soil surface (Table 10). In 2008 due to the dry weather, we were unable to collect samples to the 24" depth. Analysis of the 0-12" depth reveals that only manure injected and commercial urea surface applied and incorporated increased nitrate-N above the control (Table 16). No linear increases in nitrate-N were observed for any of the application methods.

In 2007 soil nitrate levels at the 0-12" depth measured in late July were increased above the control when manure was injected below the soil surface and when commercial fertilizer was supplied (Table 11). A linear increase in soil nitrate level was also noted with increasing N application rate for the same two treatments.

In 2007, post-harvest soil nitrate levels in the 0-12" and 12-24 depths were not significantly different from the unfertilized controls (Table 12). There were linear increases in soil nitrate levels when manure was injected and when commercial fertilizer was surface applied followed by incorporation at both depths. In 2008, post-harvest soil nitrate levels in the 0-12" depth were significantly higher than the control when manure was surface applied, regardless of tillage (Table 17). This may be an indication that the unusually cool spring delayed mineralization of some of the organic-N fraction from the manure application causing more residual nitrate-N to be present later in the year. This is corroborated by the lack of tissue N response measured during the growing season. This should not be alarming environmentally because a corn crop would have likely revealed a similar accumulation of nitrate-N.

ACKNOWLEDGEMENTS

We would like to thank the National Pork Board for their funding support of this project.

REFERENCES

- Hesterman, O.B., and T.G. Isleib. 1991. Response of first-year soybeans to row spacing, inoculation treatments, and nitrogen fertilization. *J. Prod. Agric.* 4:589-593.
- Hoitink, HAJ and M.J. Boehm. 1999. Biocontrol within the context of soil microbial communities: a substrate-dependent phenomenon. *Ann. Rev. Phytopathol.* 37:427-446.
- Hoitink, H.A.J., and A.F. Schmitthenner. 1988. Effects of composted municipal sludge on soilborne plant pathogens. Project Summary, United States Environmental Protection Agency, Water Engineering Research Laboratory, Cincinnati, OH.
- Lazarovits, G. 2001. Management of soil-borne plant pathogens with organic soil amendments: a disease control strategy salvaged from the past. *Can J. Plant Pathol.* 23:1-7.
- Osborne, S.L., and W. Riedell. 2006. Soybean growth response to low rates of nitrogen applied at planting in the northern Great Plains. *J. Plant Nutr.* (in print).
- Reese Jr., P.R., and G.R. Buss. 1992. Response of dryland soybeans to nitrogen in full-season and double crop soybeans. *J. Prod. Agric.* 5:528-531.
- Reynolds, D.A., G.L. Tylka, and C.A. Martinson. 1999. Effect of swine manure on soybean cyst nematode. National Soybean Cyst Nematode Conference, January 7-8, Orlando, FL.
- SAS Institute, Inc. 1998. *SAS/STAT User's Guide*, 6.03 Ed. SAS Institute, Inc., Cary, NC.
- Schmidt, J.P., Schmitt, M.A., Randall, G.W., Lamb, J.A., Orf, J.H. and H.T. Gollany. 2000. Swine manure application to nodulating and nonnodulating soybean. *Agron. J.* 92: 987-992.
- Schmidt, J.P., Lamb, J.A., Schmitt, M.A., Randall, G.W., Orf, J.H. and H.T. Gollany. 2001. Soybean varietal response to liquid swine manure application. *Agron. J.* 93:358-363.

Table 1. Initial manure test values, sampled winter 2005.

Moisture (%)	Total N	Ammonia-N	Organic-N	P ₂ O ₅	K ₂ O
-----lbs/1000 gal-----					
95.9	42.1	29.9	12.2	21.3	25.1

Table 2. Manure analysis as applied.

Year	Moisture (%)	Total N	Ammonia-N	Organic-N	P ₂ O ₅	K ₂ O
-----lbs/1000 gal-----						
2007	96.7	26.1	22.7	3.3	15.9	19.4
2008	92.1	41.8	18.2	23.6	32.9	19.2

Table 3. Initial soil test information collected in the fall prior to experiment establishment.

Year	Soil pH ^a	Available P ^b (mg/kg)	Exchangeable K ^c (mg/kg)	Organic matter ^d (%)
2006	5.9	17	87	2.5
2007	6.7	11	73	2.1

a-1:1 soil:water; b-Bray-Kurtz P-1; c-ammonium acetate; d-loss on ignition

Table 4. Simple effects of manure and commercial fertilizer applied on soybean dry weight, tissue N concentration, and N uptake measured on June 28, 2007.

Application method	N source	N rate (lb/acre)	Dry weight (lb/acre)	Tissue N (%)	N uptake (lb/acre)
Surface (tilled)	None	0	2940	3.5	103
Surface (no-tilled)	None	0	4029	3.5	139
Injection (tilled)	None	0	2505	3.5	87
Injection (tilled)	Manure	60	3702	3.5	128
		120	3812	3.7	140
		180	4356	3.8	168
Surface (tilled)	Manure	60	3158	3.5	109
		120	3812	3.8	135
		180	3158	3.5	113
Surface (no-tilled)	Manure	60	3376	3.7	126
		120	3158	3.6	113
		180	3812	3.4	130
Surface (tilled)	Commercial	60	3703	3.4	125
		120	2614	3.6	91
		180	4356	3.8	167
Contrasts					
			**	NS	**
			**	**	***
			NS	NS	NS
			NS	NS	NS
			NS	NS	NS
			NS	NS	NS
			NS	NS	NS
			NS	**	*

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 5. Simple effects of manure and commercial fertilizer applied on soybean dry weight, tissue N concentration, and N uptake measured on July 18, 2007.

Application method	N source	N rate (lb/acre)	Dry weight (lb/acre)	Tissue N (%)	N uptake (lb/acre)
Surface (tilled)	None	0	7296	4.3	313
Surface (no-tilled)	None	0	10890	3.9	421
Injection (tilled)	None	0	10237	4.0	411
Injection (tilled)	Manure	60	9909	4.1	411
		120	12033	4.4	528
		180	12142	4.7	565
Surface (tilled)	Manure	60	12197	4.3	530
		120	11652	4.3	505
		180	12632	4.5	564
Surface (no-tilled)	Manure	60	10237	4.1	420
		120	12088	4.3	522
		180	12088	4.5	539
Surface (tilled)	Commercial	60	11326	4.3	482
		120	16444	4.5	739
		180	16771	4.3	712
Contrasts					
			NS	**	***
			*	***	***
			***	NS	***
			***	NS	***
			NS	**	NS
			NS	***	**
			***	NS	***
			***	NS	***

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 6. Effect of manure and commercial fertilizer applied on soybean yield, 2007.

Application method	N source	N rate (lb/acre)	Soybean yield (bu/acre)
Surface (tilled)	None	0	69
Surface (no-tilled)	None	0	66
Injection (tilled)	None	0	56
Injection (tilled)	Manure	60	65
		120	75
		180	83
Surface (tilled)	Manure	60	77
		120	77
		180	77
Surface (no-tilled)	Manure	60	74
		120	80
		180	72
Surface (tilled)	Commercial	60	80
		120	79
		180	75
Contrasts			
			<i>Manure injected versus control</i> ***
			<i>Manure injected linear</i> ***
			<i>Manure surface tilled versus control</i> *
			<i>Manure surface tilled linear</i> NS
			<i>Manure surface no-till versus control</i> **
			<i>Manure surface no-till linear</i> NS
			<i>Commercial surface till versus control</i> **
			<i>Commercial surface till linear</i> NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 7. Effect of manure and commercial fertilizer applied on tap root weight, total root weight, and nematode counts at first sampling date – June 28, 2007.

Application method	N source	N rate (lb/acre)	Tap root weight (g)	Total root weight (g)	Nematode counts (/0.5g)
Surface (tilled)	None	0	1.08	1.96	15.8
Surface (no-tilled)	None	0	1.26	2.22	12.8
Injection (tilled)	None	0	1.27	2.50	12.8
Injection (tilled)	Manure	60	1.08	2.28	7.0
		120	1.09	2.81	4.3
		180	0.89	2.22	6.9
Surface (tilled)	Manure	60	1.94	3.58	3.3
		120	1.44	2.68	1.0
		180	1.13	2.17	3.5
Surface (no-tilled)	Manure	60	1.50	2.91	9.5
		120	0.98	2.14	7.5
		180	1.12	2.33	1.8
Surface (tilled)	Commercial	60	2.38	3.63	4.8
		120	2.39	4.09	12.5
		180	3.08	5.05	6.0
Contrasts					
<i>Manure injected versus control</i>			NS	NS	NS
<i>Manure injected linear</i>			NS	NS	NS
<i>Manure surface tilled versus control</i>			NS	NS	**
<i>Manure surface tilled linear</i>			NS	NS	*
<i>Manure surface no-till versus control</i>			NS	NS	NS
<i>Manure surface no-till linear</i>			NS	NS	NS
<i>Commercial surface till versus control</i>			**	***	NS
<i>Commercial surface till linear</i>			***	***	NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 8. Effect of manure and commercial fertilizer applied on tap root weight, total root weight, and nematode counts at first sampling date – July 31, 2007.

Application method	N source	N rate (lb/acre)	Tap root weight (g)	Total root weight (g)	Nematode counts (/0.5g)	
Surface (tilled)	None	0	3.6	10.7	42.3	
Surface (no-tilled)	None	0	3.1	10.2	16.3	
Injection (tilled)	None	0	3.6	9.2	15.0	
Injection (tilled)	Manure	60	2.7	8.4	6.6	
		120	3.4	7.0	15.9	
		180	2.9	8.4	14.5	
Surface (tilled)	Manure	60	1.9	7.1	1.8	
		120	2.7	9.0	4.0	
		180	3.8	9.7	41.3	
Surface (no-tilled)	Manure	60	3.5	7.2	13.8	
		120	2.4	6.5	8.8	
		180	3.6	7.6	7.5	
Surface (tilled)	Commercial	60	2.3	7.5	4.8	
		120	4.2	13.3	9.3	
		180	4.7	12.2	9.8	
Contrasts						
			<i>Manure injected versus control</i>	NS	NS	NS
			<i>Manure injected linear</i>	NS	NS	NS
			<i>Manure surface tilled versus control</i>	NS	NS	***
			<i>Manure surface tilled linear</i>	NS	NS	NS
			<i>Manure surface no-till versus control</i>	NS	NS	NS
			<i>Manure surface no-till linear</i>	NS	NS	NS
			<i>Commercial surface till versus control</i>	NS	NS	***
			<i>Commercial surface till linear</i>	NS	NS	***

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 9. Effects of manure and commercial fertilizer applied on soil nitrate concentrations at two depths (0-12” and 12-24”) measured on June 1, 2007.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration	
			0-12”	12-24”
-----mg/kg-----				
Surface (tilled)	None	0	9.1	6.5
Surface (no-tilled)	None	0	10.9	6.3
Injection (tilled)	None	0	15.0	7.5
Injection (tilled)	Manure	60	13.1	8.3
		120	27.6	8.3
		180	22.8	8.0
Surface (tilled)	Manure	60	13.6	7.0
		120	20.1	7.7
		180	21.8	8.4
Surface (no-tilled)	Manure	60	16.1	7.1
		120	17.8	7.6
		180	23.0	10.4
Surface (tilled)	Commercial	60	16.0	7.1
		120	25.9	9.6
		180	29.1	9.2
Contrasts				
			NS	NS
			***	NS
			**	NS
			***	NS
			**	**
			***	***
			***	**
			***	***

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 10. Effects of manure and commercial fertilizer applied on soil nitrate concentrations at two depths (0-12” and 12-24”) measured on June 28, 2007.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration	
			0-12”	12-24”
-----mg/kg-----				
Surface (tilled)	None	0	10.6	6.4
Surface (no-tilled)	None	0	6.4	5.2
Injection (tilled)	None	0	7.4	6.4
Injection (tilled)	Manure	60	7.9	6.3
		120	9.6	6.6
		180	23.4	9.8
Surface (tilled)	Manure	60	8.9	5.6
		120	15.4	10.1
		180	14.1	7.9
Surface (no-tilled)	Manure	60	10.8	6.2
		120	15.1	7.4
		180	19.0	8.0
Surface (tilled)	Commercial	60	10.7	6.1
		120	14.1	6.7
		180	17.5	8.3
Contrasts				
			**	NS
			***	**
			NS	NS
			NS	NS
			***	NS
			***	NS
			NS	NS
			**	NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 11. Effect of manure and commercial fertilizer applied on soil nitrate concentrations at one depths (0-12”) measured on July 18, 2007.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration (mg/kg)
Surface (tilled)	None	0	5.9
Surface (no-tilled)	None	0	5.7
Injection (tilled)	None	0	5.4
Injection (tilled)	Manure	60	7.6
		120	11.2
		180	23.3
Surface (tilled)	Manure	60	6.5
		120	7.9
		180	11.6
Surface (no-tilled)	Manure	60	6.2
		120	8.7
		180	9.1
Surface (tilled)	Commercial	60	9.0
		120	12.1
		180	19.3
Contrasts			
			<i>Manure injected versus control</i> ***
			<i>Manure injected linear</i> ***
			<i>Manure surface tilled versus control</i> NS
			<i>Manure surface tilled linear</i> NS
			<i>Manure surface no-till versus control</i> NS
			<i>Manure surface no-till linear</i> NS
			<i>Commercial surface till versus control</i> **
			<i>Commercial surface till linear</i> ***

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 12. Effect of manure and commercial fertilizer applied on soil nitrate concentrations at two depths (0-12” and 12-24”) measured in September 25, 2007.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration	
			0-12”	12-24”
-----mg/kg-----				
Surface (tilled)	None	0	7.4	4.2
Surface (no-tilled)	None	0	7.7	4.5
Injection (tilled)	None	0	6.7	4.1
Injection (tilled)	Manure	60	6.7	4.1
		120	7.4	5.4
		180	11.4	6.3
Surface (tilled)	Manure	60	8.2	4.6
		120	8.4	5.2
		180	6.9	5.0
Surface (no-tilled)	Manure	60	7.3	4.3
		120	7.1	5.0
		180	8.9	4.6
Surface (tilled)	Commercial	60	7.3	3.5
		120	11.1	4.2
		180	10.0	6.0
Contrasts				
			NS	NS
			**	***
			NS	NS
			*	*

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 13. Simple effects of manure and commercial fertilizer applied on soybean dry weight, tissue N concentration, and N uptake at V4 measured on July 3, 2008.

Application method	N source	N rate (lb/acre)	Dry weight (lb/acre)	Tissue N (%)	N uptake (lb/acre)
Surface (tilled)	None	0	987	3.5	34.1
Surface (no-tilled)	None	0	987	3.6	35.8
Injection (tilled)	None	0	953	3.6	34.2
Injection (tilled)	Manure	60	953	3.5	33.4
		120	1123	3.4	38.6
		180	1157	3.8	43.6
Surface (tilled)	Manure	60	1089	3.6	39.6
		120	987	3.7	36.4
		180	1225	3.8	46.7
Surface (no-tilled)	Manure	60	1123	3.6	40.8
		120	1021	3.5	35.6
		180	953	3.8	36.0
Surface (tilled)	Commercial	60	1395	3.9	54.9
		120	1259	3.9	48.9
		180	1259	4.0	50.3
Surface (no-tilled)	Commercial	60	1055	3.6	38.0
		120	953	3.6	34.9
		180	1021	3.6	36.2
Contrasts					
			NS	NS	NS
			*	NS	*
			NS	*	NS
			NS	*	**
			NS	NS	NS
			NS	NS	NS
			***	**	***
			NS	**	**
			NS	NS	NS
			NS	NS	NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 14. Simple effects of manure and commercial fertilizer applied on soybean dry weight, tissue N concentration, and N uptake at R1 measured on August 7, 2008.

Application method	N source	N rate (lb/acre)	Dry weight (lb/acre)	Tissue N (%)	N uptake (lb/acre)
Surface (tilled)	None	0	3921	3.4	134.0
Surface (no-tilled)	None	0	3757	3.3	126.0
Injection (tilled)	None	0	4710	3.5	162.7
Injection (tilled)	Manure	60	3430	3.3	112.5
		120	4765	3.7	175.3
		180	5064	3.5	176.3
Surface (tilled)	Manure	60	4765	3.7	174.1
		120	4520	3.6	161.9
		180	6480	3.6	231.4
Surface (no-tilled)	Manure	60	3757	3.9	144.6
		120	4656	3.7	173.1
		180	5500	3.7	204.7
Surface (tilled)	Commercial	60	4737	3.3	159.4
		120	5009	3.6	179.1
		180	5636	3.6	197.6
Surface (no-tilled)	Commercial	60	5009	3.5	177.0
		120	3648	3.8	138.6
		180	4519	3.5	157.5
Contrasts					
			NS	NS	NS
			NS	NS	NS
			**	*	***
			***	NS	***
			NS	***	**
			**	**	***
			**	NS	**
			**	**	***
			NS	NS	NS
			NS	NS	NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 15. Effects of manure and commercial fertilizer applied on soil nitrate concentrations at two depths (0-12” and 12-24”) measured on July 3, 2008.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration		
			0-12”	12-24”	
-----mg/kg-----					
Surface (tilled)	None	0	8.9	5.6	
Surface (no-tilled)	None	0	7.9	4.8	
Injection (tilled)	None	0	9.7	6.7	
Injection (tilled)	Manure	60	10.4	5.8	
		120	12.2	8.5	
		180	17.2	6.9	
Surface (tilled)	Manure	60	13.3	7.5	
		120	13.1	6.2	
		180	18.4	10.1	
Surface (no-tilled)	Manure	60	9.1	5.9	
		120	11.5	7.5	
		180	15.3	6.7	
Surface (tilled)	Commercial	60	20.7	9.6	
		120	22.1	11.6	
		180	24.9	12.9	
Surface (no-tilled)	Commercial	60	6.3	4.3	
		120	7.4	5.4	
		180	8.9	5.7	
Contrasts					
			<i>Manure injected versus control</i>	NS	NS
			<i>Manure injected linear</i>	**	NS
			<i>Manure surface tilled versus control</i>	**	*
			<i>Manure surface tilled linear</i>	***	**
			<i>Manure surface no-till versus control</i>	NS	NS
			<i>Manure surface no-till linear</i>	**	NS
			<i>Commercial surface till versus control</i>	***	***
			<i>Commercial surface till linear</i>	***	***
			<i>Commercial surface no-till versus control</i>	NS	NS
			<i>Commercial surface no-till linear</i>	NS	NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 16. Effects of manure and commercial fertilizer applied on soil ammonium and nitrate concentrations at one depth (0-12”) measured on August 7, 2008.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration 0-12” -----mg/kg-----
Surface (tilled)	None	0	4.4
Surface (no-tilled)	None	0	4.3
Injection (tilled)	None	0	4.3
Injection (tilled)	Manure	60	6.1
		120	6.5
		180	5.5
Surface (tilled)	Manure	60	5.1
		120	4.6
		180	5.4
Surface (no-tilled)	Manure	60	5.1
		120	5.8
		180	4.9
Surface (tilled)	Commercial	60	5.7
		120	6.6
		180	5.4
Surface (no-tilled)	Commercial	60	4.2
		120	4.4
		180	4.4
Contrasts			
<i>Manure injected versus control</i>			**
<i>Manure injected linear</i>			NS
<i>Manure surface tilled versus control</i>			NS
<i>Manure surface tilled linear</i>			NS
<i>Manure surface no-till versus control</i>			NS
<i>Manure surface no-till linear</i>			NS
<i>Commercial surface till versus control</i>			*
<i>Commercial surface till linear</i>			NS
<i>Commercial surface no-till versus control</i>			NS
<i>Commercial surface no-till linear</i>			NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 17. Effects of manure and commercial fertilizer applied on soil nitrate concentrations at one depth (0-12”) measured on October 30, 2008.

Application method	N source	N rate (lb/acre)	Nitrate-N concentration 0-12”
Surface (tilled)	None	0	5.5
Surface (no-tilled)	None	0	5.7
Injection (tilled)	None	0	5.7
Injection (tilled)	Manure	60	5.0
		120	6.6
		180	7.0
Surface (tilled)	Manure	60	7.4
		120	9.5
		180	9.5
Surface (no-tilled)	Manure	60	7.4
		120	9.9
		180	9.7
Surface (tilled)	Commercial	60	5.5
		120	6.1
		180	6.4
Surface (no-tilled)	Commercial	60	5.4
		120	5.5
		180	5.5
Contrasts			
	<i>Manure injected versus control</i>		NS
	<i>Manure injected linear</i>		NS
	<i>Manure surface tilled versus control</i>		***
	<i>Manure surface tilled linear</i>		***
	<i>Manure surface no-till versus control</i>		***
	<i>Manure surface no-till linear</i>		***
	<i>Commercial surface till versus control</i>		NS
	<i>Commercial surface till linear</i>		NS
	<i>Commercial surface no-till versus control</i>		NS
	<i>Commercial surface no-till linear</i>		NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;

Table 18. Effect of manure and commercial fertilizer applied on soybean yield, 2008.

Application method	N source	N rate (lb/acre)	Soybean yield (bu/acre)
Surface (tilled)	None	0	38
Surface (no-tilled)	None	0	36
Injection (tilled)	None	0	40
Injection (tilled)	Manure	60	31
		120	46
		180	46
Surface (tilled)	Manure	60	49
		120	51
		180	59
Surface (no-tilled)	Manure	60	47
		120	53
		180	56
Surface (tilled)	Commercial	60	53
		120	49
		180	51
Surface (no-tilled)	Commercial	60	43
		120	43
		180	44
Contrasts			
		<i>Manure injected versus control</i>	NS
		<i>Manure injected linear</i>	**
		<i>Manure surface tilled versus control</i>	***
		<i>Manure surface tilled linear</i>	***
		<i>Manure surface no-till versus control</i>	***
		<i>Manure surface no-till linear</i>	***
		<i>Commercial surface till versus control</i>	***
		<i>Commercial surface till linear</i>	***
		<i>Commercial surface no-till versus control</i>	NS
		<i>Commercial surface no-till linear</i>	NS

***, **, * - significant at the 0.01, 0.05, 0.1 probability levels, respectively; NS – non-significant;