

Title: A review of heavy weight market pigs: status of knowledge and future needs assessment – **NPB #15-203**

Investigator: J. M. DeRouchey*²

Co-Investigators: F. Wu,* K. R. Vierck,* T. G. O'Quinn,* M. D. Tokach,* R. D. Goodband,* S. S. Dritz,[†] and J. C. Woodworth*

Institution: *Department of Animal Science, Kansas State University, Manhattan 66506; and [†]Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University, Manhattan 66506

Date Submitted: May 31, 2016

¹ Funding, wholly or in part, was provided by *The National Pork Board*. The authors acknowledge A. D. Santos and D. A. Marçal for their assistance.

² Corresponding author: jderouch@ksu.edu

These research results were submitted in fulfillment of checkoff-funded research projects. This report is published directly as submitted by the project's principal investigator. This report has not been peer-reviewed.

For more information contact:

National Pork Board • PO Box 9114 • Des Moines, IA 50306 USA • 800-456-7675 • Fax: 515-223-2646 • pork.org

ABSTRACT: Marketing weight is an important economic variable that impacts the productivity and profitability of finishing pig production. Marketing weight has been increasing worldwide over the past decades driven by the dilution of fixed production cost over more weight per pig and the improvement of genetic selection of lean-type pigs. This review was aimed to summarize current knowledge and assess the future research needs on producing finishing pigs with marketing weight greater than 130 kg. Based on a thorough literature review, increasing marketing weight affected overall pig growth; in particular, cumulative ADG decreased by 4.0 g/d, ADFI increased by 78.1 g/d, and G:F decreased by 0.011 for every 10 kg increase of marketing weight. Increasing marketing weight by 10 kg increased carcass yield by 0.41 percentage units, backfat by 1.8 mm, LM area by 1.8 cm², carcass length by 2.2 cm, and belly yield by 0.32 percentage units, but decreased percentage of fat-free-lean by 0.78 units and loin, shoulder, and ham yields by 0.13, 0.16, and 0.17 percentage units, respectively. Studies that investigated the effects of marketing weight on pork quality observed decreased pH by 0.02 and 0.01 at 45 min and 24 h postmortem, respectively, and increased a* value by 0.28 per 10 kg marketing weight increase. Heavier market pigs had increased concentrations of saturated fatty acids and intramuscular fat. However, studies reported conflicting results for L* and b* values, drip loss, Warner-Bratzler shear force, and sensory properties of pigs in response to increasing marketing weight. A limited amount of research has been conducted to provide nutritional requirement guideline for pigs greater than 140 kg. Increased weight and size of heavy pigs can create challenges to farm and packer facilities and equipment. Discussions and recommendations are provided concerning the adjustments for floor and feeder space, barn design, ventilation, disease control, transportation, and carcass processing needed for increasing marketing weight. In conclusion, increasing marketing weight creates both opportunities and challenges to current finishing pig production,

and future research is needed to provide nutritional and management guidelines and improve feed efficiency and meat quality of heavy weight market pigs.

Key words: carcass quality, growth, heavy pig, marketing weight, meat quality

INTRODUCTION

Marketing weight is an important variable that affects the profitability of finishing pig production due to its impact on pig growth efficiency and the quantity and quality of pork produced. Average marketing weight in the U.S. has been steadily increasing for over 80 years and increased from 121.1 kg in 2004 to 125.6 kg in 2013 (NASS, 2014). A dilution of fixed production cost is a major force that drives the increase of marketing weight because the total number of pigs required to produce a given quantity of pork is reduced (Park and Lee, 2011). A drawback of the increased marketing weight is reduced G:F resulting from accelerated fat accretion and a declining rate of lean deposition during in the late finishing phase (Shields et al., 1983; Gu et al., 1991; Piao et al., 2004). In addition, increased weight and size of heavy pigs creates challenges to farm facilities and equipment, such as floor and feeder space, ventilation, and transportation systems, which in turn influences pig growth performance.

Some additional factors that require consideration when increasing marketing weight include genetic selection and nutritional requirements. Lean-genotype pigs are needed to prolong the period of efficient weight gain, while the selection for lean gain rate should also be balanced with the requirements of meat quality and health attributes. From a nutritional prospective, nutrient requirements are established for pigs less than 140 kg (NRC, 2012); however, pigs with further increased BW have greater maintenance needs than lighter BW and, therefore, require additional research to provide nutritional guidelines. Finally, information regarding the impact of meat quality with increasing marketing weight, such as color, primal cut yields, and intramuscular fatness of heavy pigs and its subsequent impact on consumer preference are needed. This review aimed to evaluate published studies involving genetic selection, nutritional requirements, health, welfare, and pork quality of heavy weight market pigs and to assess the future research needs.

MATERIALS AND METHODS

Examination of published studies was conducted via the Kansas State University Libraries, using databases including AGRICOLA, CAB International, MEDLINE, National Pork Board Research Database, and SCOPUS. No year of publication limits was set in any of the electronic database searches. Additional search of literature was performed within the following journals: *Journal of Animal Science*, *Animal*, *Animal Feed Science and Technology*, *Meat Science*, *Livestock Science*, and *Livestock Production Science*. Key words used for the above databases included: “heavy pig*”, “heavy hog*”, “heavy weight”, “finishing pig*”, “finishing hog*”, “late finishing pig*”, “late finishing hog*”, “slaughter weight”, “harvest weight”, “marketing weight”, along with the key words associated with the aspects of selection/genetics, nutrition, pork/meat quality, pork safety, and swine health and well-being. In addition, non-peer-reviewed publications (i.e., university extension and company reports) were also collected, closely scrutinized for accuracy and quality, and served as valuable resources of information for this review. Conference proceedings and abstracts that were not included in the peer-reviewed databases were searched using Searchable Proceedings of Animal Conferences (S-PAC) and Google Scholar search engine. Additionally, personal communication with genetic and production companies, university researchers, and packing plant personnel were performed for the collection of internally-generated information that had application for this review.

In this review, heavy weight market pigs refer to pigs with marketing weight greater than 130 kg. For the summary of the effects of marketing weight on pig growth performance, carcass characteristics, and pork quality, the data set excluded studies in which the greatest marketing weight used was less than 125 kg and pigs were restricted fed during the experiment. Sensitivities of growth, carcass, and pork quality traits in response to increasing marketing weight by 10 kg were generated using simple linear regression. These analyses were based on the

assumption that traits had linear responses to the increase in marketing weight and there were no interactive effects between marketing weight and other factors (i.e., gender, inclusion of growth promoters). Average responses were reported as the mean among studies. In the calculation of average responses, studies by Latorre et al. (2004 and 2008) were excluded for ADG, ADFI, and G:F, because pigs were reported to be under heat stress, and a study by Serrano et al. (2008) was excluded for growth and carcass traits due to the use of Iberian obese pig breed that is typically not used in north America pig production. A study by Piao et al. (2004) was excluded in the calculation for drip loss due to the abnormally high value reported (greater than 3 standard deviations from the mean of all values).

RESULTS AND DISCUSSIONS

Impact of marketing weight on growth performance, carcass characteristics, and meat quality

Growth performance

Numerous studies have been conducted to investigate the effects of increasing marketing weight on growth performance (i.e., ADG, ADFI, and G:F) of growing-finishing or finishing pigs. A total of 14 experiments involving pigs harvested at weights greater than 125 kg were summarized in Table 1. Although instantaneous gain rate and feed intake of pigs follow allometric patterns as BW increases (sigmoid growth curve; Schinckel et al., 2006; Shull, 2013), we plotted the cumulative ADG and ADFI values against marketing weight reported by the reviewed studies and observed linear growth responses to increasing marketing weight. Eight out of the 14 reviewed studies reported a decrease in cumulative ADG of 3.6 to 54.9 g/d for every 10 kg increase in marketing weight, whereas the remaining studies showed increased ADG of 2.8 to 8.7 g/d when marketing weight increased by 10 kg. Cumulative ADFI was reported in 13 studies with ADFI increasing by 52.7 to 163.6 g/d in 11 studies. Conversely, ADFI decreased by 3.0 and 78.0

g/d in 2 studies (Latorre et al., 2004 and 2008, respectively) where heat stress of pigs under severe summer weather was reported. Reduction in cumulative G:F was observed in all the reviewed studies with the magnitude varying from 0.003 to 0.017 units per 10 kg marketing weight increase. On average (calculation excluded data from Latorre et al., 2004 and 2008 due to suppressed ADFI and data from Serrano et al., 2008 due to the use of an Italian obese pig breed), increasing marketing weight by 10 kg decreased cumulative ADG by 4.0 g/d, increased ADFI by 78.1 g/d, and decreased G:F by 0.011.

It is not surprising that pigs marketed at heavy weights have elevated ADFI, because the increased body size and physical capacity of the digestive tract improve the ability of pigs to consume more feed (Suarez-Belloch et al., 2013). Nevertheless, the efficiency of BW gain declines greatly during the late growth stages, which is attributed to accelerated fat accretion and declining rates of water and protein deposition (Shields et al., 1983; Gu et al., 1991; Piao et al., 2004). Increased maintenance requirements in heavy finishing pigs may also contribute to decreased G:F (Gu et al., 1991). For the ADG response, researchers (Schinckel et al., 2006; Jungst et al., 2012a,b; Shull, 2013) have demonstrated that the instantaneous growth rate of growing-finishing pigs (average between barrow and gilt) reaches a plateau at an average BW of 78 to 85 kg and decreases thereafter. However, evaluating data from the 14 experiments, it is hard to accurately describe reason(s) why cumulative ADG was improved in half of the experiments and diminished in the other half. Possible explanations of this discrepancy can be proposed. First, nutritional programs used and, particularly, the dietary energy and protein supply, varied among these studies, which would influence the growth responses of pigs at increasing marketing weight. Secondly, selection of the initial and terminal BW differed among studies and could also be a factor. Generally, the greater the initial BW (shorter overall feeding period) and wider range of marketing weight used, the more prominent responses were observed. However, use of wide

marketing weight range tended to result in a quadratic response of cumulative ADG to increasing marketing weight (Shull, 2013), which also affected the precision of linear quantification for ADG. Thirdly, housing system and especially the floor and feeder spaces allowance could affect the ADFI and, subsequently, ADG of pigs. Furthermore, dissimilar genetic lines of pigs used in the studies had varied growth patterns at heavy weights. Lean-type pigs are desired for producing pigs marketed at heavy weights (Kim et al., 2005). However, some of the reviewed studies were carried out on pigs that were aimed for dry-cured ham production (Latorre et al., 2004 and 2008; Serrano et al., 2008), which were often selected for high fat thickness; discrepant growth responses of these pigs could be expected when compared with modern lean-type pigs. Finally, quantification of growth responses is also determined by the methodology used in the studies. Only studies reporting cumulative growth responses were compared herein because relatively few studies (Carr et al., 1978; Gu et al., 1991; Shull, 2013) in the literature reported instantaneous growth rate and the methodologies utilized to measure instantaneous growth rate differed among these studies.

Carcass characteristics

Increasing marketing weight greatly influences carcass characteristics of pigs. For this analysis, 25 studies were reviewed where carcass traits of pigs with increasing market weight were determined (Table 2). Twenty studies evaluated the percentage carcass yield of pigs harvested at heavy weights; increased yield was documented in 19 studies ranging from 0.05 to 1.05 percentage units per 10 kg increase in marketing weight. Across all studies, the mean increase in carcass yield was approximately 0.41 percentage units per 10 kg marketing weight increase. Increased carcass yield was due to a greater allometric coefficient of carcass than the whole body (Gu et al., 1992). Shields et al. (1983) suggested that the carcass only represented 70% of the live weight at 56 kg, but 79% by 146 kg; whereas the relative proportion of the

intestinal tract decreased from 5.6% to 4.3%, and that of internal organs also decreased from 4.5% to 3.2%. However, one study reported a reduced yield of 0.49 percentage units per 10 kg increase of marketing weight (Piao et al., 2004).

All studies considered in this review observed an increase in backfat thickness with increased marketing weight. However, increases in backfat varied among studies, ranging from 0.5 to 3.0 mm per 10 kg marketing weight increase. Across the studies reviewed, there was an average increase in backfat of 1.8 mm per 10 kg increase in marketing weight. In terms of overall fat deposition, there is little published research evaluating specific areas of deposition, with the exception of the belly and back fat. Correa et al. (2008) reported significant increases in belly fat thickness as marketing weight increased from 107 to 125 kg, though no other studies have evaluated this trait.

Percentage fat-free lean was reduced in most cited studies with increased marketing weight. The observed reduction in percentage fat-free lean was most likely due to the increased backfat found in heavy pigs. In contrast, 3 studies found an increase in percentage fat-free lean ranging from 0.05 to 2.28 unit per 10 kg increase in marketing weight. Interestingly, the studies reported an increase in percentage fat-free lean were those that used greater initial BW and narrow ranges between initial and marketing weights than other studies.

As marketing weight increases, there is a general trend of increasing LM area and carcass length, which can be explained by the greater body size of heavy pigs. Almost all the reviewed studies found an increase in LM area ranging from 1.1 to 2.7 cm², with an average of 1.95 cm² per 10 kg marketing weight increase. Conversely, Leach et al. (1996) found a 0.02 cm² decrease per 10 kg increase in body weight when comparing the effect of the halothane gene on pigs with increasing BW from 110 to 140 kg. All the 25 studies observed increasing carcass length with greater marketing weights. However, wide variation of the increase in carcass length was present

ranging from 1.3 to 3.1 cm, with an average of 2.2 cm, per 10 kg of additional BW. Increased carcass length may cause issues in processing plants if pigs are too large to fit through typical equipment, such as rails, scalders, carcass splitters, and other mechanized fabrication equipment.

A total of 14 studies evaluated the effects of increasing marketing weight on subprimal cut yields. Belly yield increased with increasing marketing weight in all studies, ranging from only 0.09 to 0.61 percentage units per 10 kg marketing weight increase. In regards to lean primal cuts, yields were generally decreased. Ten studies observed decreased loin yield, ranging from 0.09 to 0.38 percentage units per 10 kg marketing weight increase. However, Cisneros et al. (1996) reported an increase in loin yield of 0.4 percentage yield per 10 kg increase in marketing weight. Of the 10 studies that evaluated shoulder yield, 7 studies reported a decrease ranging from 0.48 to 0.02 percentage units per 10 kg marketing weight increase. However, 3 studies found a slight increase in shoulder yield ranging from 0.08 to 0.09 percentage units per 10 kg marketing weight increase. Ham yield was affected similarly to shoulder and loin yields. As marketing weight increased, ham yield decreased in 10 out of the 13 studies. Decreases in ham yield ranged from 0.09 to 0.36 percentage units per 10 kg increase in marketing weight. However, 3 studies reported slight increases in ham yield; this might be related to how the loin was removed, as Latorre et al. (2004) and Serrano et al. (2008) were studies done with Italian heavy weight pigs. In addition, it is important to note that changes of primal cut yields were influenced by whether the data reported trimmed or untrimmed cuts. More prominent responses could be expected for untrimmed cuts because a great amount of fat was deposited on the cuts during the last stages of growth. On average, increasing marketing weight by 10 kg increased belly yield by 0.32 percentage units, but reduced loin, shoulder, and ham yields by 0.13, 0.16, and 0.17 percentage units, respectively.

Pork Quality

Pork quality is important for several reasons, including product functionality, consumer preference, and palatability. Several studies have evaluated pork quality traits as it relates to increased marketing weight. These include: pH, drip loss, cooking loss, Warner-Bratzler shear force, intramuscular fat or marbling scores, iodine value, as well as instrumental color scores and sensory panel data.

The majority of published literature has observed a decrease in pH as carcass weight increases. Decreased pH negatively affects drip loss, color, and several other pork quality traits. All the 6 studies reported initial pH measured at 45 min to 1 h postmortem, and 6 out of 8 studies evaluated ultimate pH at 24 h postmortem observed decreased pH values when increasing marketing weights. Beattie et al. (1999) and Martin et al. (1980) found significant decreases in pH at 1 h postmortem, but no significant differences at 24 h or in ultimate pH when comparing pigs with increasing marketing weight from 92 to 131 kg and 73 to 137 kg, respectively. Additionally, Martin et al. (1980) also reported a negative, but weak, correlation ($r = -0.05$) between carcass weight and 1 h pH. When comparing pigs at 8 months of age with 143.6 kg BW versus those 10 months of age with 181.8 kg BW, Virgili et al. (2003) observed a 0.05 unit reduction in pH of the semimembranosus at 1 h as well as at 24 h as marketing weight increased by 10 kg. Moreover, Cisneros et al. (1996) reported a reduction of pH at a rate of 0.01 unit at 45 min and a 0.02 unit reduction at 24 h postmortem per 10 kg of additional BW. Park and Lee (2011) observed a 0.02 unit reduction in 24 h pH per 10 kg increase in marketing weight from 116 to 133 kg. In a study involving pigs with increasing marketing weight from 120 to 170 kg, Durkin et al. (2012) observed a quadratic response of pH at 45 min postmortem. In that study, pH of semimembranosus increased by 0.01 unit per 10 kg increase in marketing weight from 120 to 140 kg and decreased at a similar rate when marketing weight increased from 140 to 170 kg. In

contrast, Piao et al. (2004) and Bertol et al. (2015) observed increases in ultimate pH at 0.02 and 0.01 respectively per 10 kg marketing weight increase.

With a reduction in pH, especially at 24 h, other pork quality factors, specifically instrumental color and drip loss are affected. Color is the number one factor affecting consumer decisions when purchasing meat, as it is used as an indicator of freshness (Mancini and Hunt, 2005). In regards to color, there are conflicting results related to increased market weights.

Overall, 8 studies have evaluated instrumental color in heavy weight carcasses. An example of the conflicting results can be found with L*, an instrumental color measurement used to evaluate the lightness or darkness of a product (greater L* value indicates a lighter color). Durkin et al. (2012) observed no significant differences in L* when comparing 120, 130, 140, 150, 160 kg pigs to those weighing greater than 170 kg. Park and Lee (2011) also observed no significant differences in L* values among pigs weighing 116, 124, and 135 kg. In contrast, Latorre et al. (2004) found a 2.48 unit reduction in L* value with a 10 kg increase in marketing weight when comparing pigs from 116 to 133 kg. In addition, when evaluating differences among pigs slaughtered at 144 and 182 kg, Virgili et al. (2003) determined a 0.01 unit reduction in L* value in the semimembranosus with every 10 kg increase in BW.

In the 7 studies evaluating a* value, an instrumental color measurement used to determine redness of a product (greater a* value indicates a more reddish color), most published literature found an increase or no significant differences as carcass weight increased. Increases in a* value were observed by Durkin et al. (2012) and Latorre et al. (2004). Durkin et al. (2012) found a 0.33 unit increase in CIE (Commission Internationale de l'Eclairage color system) a* values in the semimembranosus muscle when comparing pigs weighing 120, 130, 140, 150, 160 kg to those weighing greater than 170 kg. Latorre et al. (2004) observed a* value increased by 0.43 units per 10 kg marketing weight increase when evaluating the effects of gender on meat

quality of pigs weighing 116, 124, and 133 kg. However, other studies found no significant differences in a^* value with increasing carcass weights (Park and Lee, 2011; Virgili et al., 2003), thus providing no clear evidence as to the effect of increased carcass weights on a^* instrumental color values.

The evaluation of b^* is an instrumental determination of yellowness in meat (greater b^* value indicates more yellowish color). Much like L^* value, the 7 studies that evaluated meat color found contradictory findings, with 4 studies finding increased values and 3 studies finding reduced values. Durkin et al. (2012) reported an increase of 0.1 unit in b^* value per 10 kg marketing weight increase. When evaluating the differences in meat quality and carcass characteristics among 8 and 10 month old Italian pigs weighing 144 and 182 kg, respectively, Virgili et al. (2003) determined there was a 0.17 unit reduction in b^* values in the semimembranosus per 10 kg marketing weight increase. Overall as marketing weights increased, there are conflicting results on instrumental color, especially in L^* and b^* values in published literature. However, such changes in instrumental color values may be of little biological significance, but may result in a minimal impact on consumer preference.

Drip loss, a measurement of waterholding capacity, is readily affected by both pH and chilling method. Of the studies evaluating the effects of increasing carcass weight on pork quality, 9 studies evaluated drip loss with conflicting results reported. With increasing BW, drip loss was increased in 5 studies, decreased in 3 studies, and inconsistent response of drip loss to increasing marketing weight was observed in 1 study. Cisneros et al. (1996) and Park and Lee (2011) found a 0.29 percentage unit increase in drip loss per additional 10 kg of BW. In addition, Martin et al. (1980) determined that carcass weight was negatively related ($r = -0.31$) to percentage expressible juice. As age and carcass weight increased, Virgili et al. (2003) observed a 0.34 percentage unit increase in drip loss for every 10 kg increase in marketing weight from

144 to 182 kg. Durkin et al. (2012) reported that drip loss of pigs marketed at 140 kg was approximately 3% less than pigs marketed at 130, 150, and 160 kg, but was not different from those marketed at 120, 140, and 170 kg. Methodology reported by these studies did not indicate any differences in chilling methods that may have affected drip loss results.

After an intensive review of heavy weight market pig literature, there is no published research on the impact of chilling rate on meat quality traits with heavy weight market pigs. Research is needed to evaluate if increased wind speeds and decreased cooler temperatures are needed to appropriately chill heavier carcasses to prevent undesirable meat quality traits.

Pork fat quality is important for product functionality and use. Three studies have evaluated the effects of increasing carcass weight on the fatty acid profiles of pork carcasses. All of the studies observed non-significant differences in MUFA among pigs of different BW (Correa et al., 2008; Lo Fiego, et al., 2005; Raj et al., 2010). In a study by Raj et al. (2010), where pigs weighing 90, 110, and 130 kg were evaluated for subcutaneous fatty acid profiles, concentrations of PUFA were reduced by 0.37 percentage units per 10 kg marketing weight increase from 90 to 130 kg. Conversely, SFA contents were increased by 0.46 percentage units per 10 kg marketing weight increase when comparing pigs weighing 90 and 130 kg (Raj et al. 2010). When examining the fatty acid profiles of fat coverings of hams in Italian heavy pigs weighing 151, 164, and 176 kg, Lo Fiego et al. (2005) observed similar results to Raj et al. (2010); as BW increased, there was a 0.36 percentage unit increase in SFA content for every 10 kg increase in marketing weight. In addition, these authors reported significant reductions in PUFA concentration as marketing weight increased; Lo Fiego et al. (2005) reported a 0.52 percentage unit reduction and Raj et al. (2010) observed a 0.37 percentage unit reduction in PUFA concentration per 10 kg increase in marketing weight. Conversely, in a study comparing bellies from heavy weight market pigs intended for cured ham production, Correa et al. (2008)

observed a tendency ($P = 0.06$) for increased PUFA content when comparing pigs weighing 107, 115, and 125 kg. However, Lo Fiego et al. (2005) observed a 0.72 unit decrease in iodine value per 10 kg increase of marketing weight. Iodine value does not affect bellies' functionality when ranging from 70 to 75 g/100g (Benz et al., 2011). Iodine values reported by Correa et al. (2008) and Lo Fiego et al. (2005) did not exceed this acceptance range, suggesting that an increase in marketing weight resulted in minimal reductions in pork product functionality.

Warner-Bratzler shear force (WBSF) results are conflicting in studies evaluating increasing marketing weight. Of the 6 studies that evaluated WBSF, Beattie et al. (1999) and Latorre et al. (2004) observed no significant differences when comparing pigs weighing 70, 80, 90, and 100 kg, as well as 116, 124, and 133 kg, respectfully. On the contrary, Cisneros et al. (1996) observed a slight reduction of 0.08 kg per 10 kg marketing weight increase in WBSF, which may be due to the increased intramuscular fat content associated with increased carcass weights. Martin et al. (1980) also observed a slightly positive, significant relationship between increasing carcass weights and shear force ($r = 0.08$), which indicated a tougher product with increasing marketing weight. In addition, Durkin et al. (2012) reported a quadratic effect of BW on tenderness; pigs weighing 140 and 160 kg had greater WBSF values and, therefore, were more tender than those weighing 120, 150 and 170 kg.

Marbling or intramuscular fat is a primary driver for both juiciness and tenderness in pork products (Cannata et al., 2010). Multiple studies (Cisneros et al., 1996; Huff-Lonergan et al., 2002; Park and Lee, 2011) demonstrated a concurrent increase in intramuscular fat in the longissimus dorsi muscle as carcass weight increases, with an exception that Martin et al. (1980) observed a weak, negative response ($r = -0.02$) of marbling to increasing carcass weight from 73 to 137 kg.

There were only 3 studies evaluated the sensory properties of heavy weight market pigs and have produced mixed results. Huff-Lonergan et al. (2002) observed significant, positive responses of juiciness ($r = 0.09$) and off-flavor presence ($r = 0.14$) to increasing carcass weight. Increase in off-flavors is likely a result of increased PUFA concentration along with enhanced fat deposition in heavy pigs (Correa et al., 2008). Contrary to those findings, Cisneros et al. (1996) observed decreased tenderness and juiciness by 0.1 and 0.04%, respectively, for every 10 kg increase in marketing weight from 100 to 160 kg. Park and Lee (2011) observed increased presence of off-flavor in raw pork as marketing weight increased from 116 to 133 kg; however, after cooking, there were no significant differences in flavor profiles. Further research is needed to determine the true effects of increasing carcass weight on sensory panel ratings.

After a thorough literature review, it was determined that there has been no research evaluating the effects of heavy marketing weight on pork safety, such as microbiological populations, antimicrobial treatments, or the potential associated dilution of sprayed-on antimicrobials (i.e., organic acids) due to increased cut and carcass size.

Factors to consider when increasing marketing weight

Genetics

Genetic selection of pigs with high lean-gain potential is essential for the production of heavy pigs. Neely et al. (1979) observed that pigs selected from lean litters (sorted based on backfat) had slower weight gain during the early stages of growth (15 to 86 kg) but gained at a faster rate thereafter compared with pigs from fat litters. During the last finishing period, lean-type pigs have less deposition of fat, thus exhibit better feed efficiency compared with non-lean genotypes (Kim et al., 2005; Park and Lee, 2011). Growth performance and carcass traits of heavy pigs varied considerably when different genetic lines are assessed. In a study where pig growth of 5 genotypes were compared at 3 BW (100, 114, and 127 kg), Gu et al. (1991) observed

that there was no genotype \times BW interaction and the difference among genotypes could be as large as 11.0, 7.3, and 14.0% for ADG, ADFI, and G:F, respectively. Similarly, Latorre et al. (2003) compared pigs bred from 3 sire lines at 2 marketing weights (122 vs. 136 kg). There were no genotype by marketing weight interactions and differences of 3.3, 1.6, and 4.9% for ADG, ADFI, and G:F, respectively, were observed. More recently, a breeding stock company (PIC, Hendersonville, TN) evaluated 2 different genotypes (PIC280 vs. PIC359) fed to 145 kg; a 2.7 kg difference was observed between lines on final BW, driven by significant differences in ADG (18 g), ADFI (90 g), and G:F (0.006 g/g; personal communication, 2016).

Effects of genetic line on carcass characteristics should also be considered when increasing marketing weight. Using 5 genotypes and 2 marketing weights (130 and 160 kg), Peloso et al. (2010) demonstrated that different genetic backgrounds was responsible for dissimilar deposition rates of fat and lean during the transition of increasing marketing weight and led to significantly varied HCW, backfat thickness, and LM depth of pigs at harvest. Pigs from different genetic lines also exhibit varied patterns in partitioning fat towards intramuscular, subcutaneous (backfat), or internal (kidney) sites at heavy weights, which contributes to a difference in meat quality among genotypes (Franci et al., 2001).

Nutrition

In general, heavy pigs have decreased requirements for dietary protein concentration (Crovetto et al., 1999, Galassi et al., 2010), likely due to decreased lean gain compared with lighter finishing pigs. Limited information is available regarding the nutritional requirements of heavy pigs over 140 kg. The NRC (2012) growth model estimates a SID Lys requirement of 0.53% (assuming corn-soybean meal diet which would contain 2,350 kcal NE/kg) for finishing pigs with 130 kg BW, which is decreased to 0.49% at 140 kg BW. However, it is important to note that these estimates have not been validated by empirical studies. Using factorial

approaches, Manini et al. (1997) predicted that the SID Lys requirement of a 120 kg pig was 0.48%, and the value was reduced to 0.44 and 0.41% of the diet for pigs with 140 and 160 kg BW, respectively. Although the change of SID Lys requirement appears to be marginal, adjustment of diet formulation or an additional feeding phase should be considered as marketing weight increases. This is because a slight decrease in feed cost during late finishing phase can be economically significant due to the high ADFI of heavy pigs. In addition, tissue turnover rates and maintenance requirements change as the pig grows, the ideal AA to Lys ratios may change with pig weight (Mahan et al., 1998a). For example, Thr, Met, and Trp are needed in greater concentrations relative to Lys in older than in younger pigs (Hahn and Baker, 1995), possibly due to a greater requirement for maintenance than for growth purposes. Furthermore, dietary P requirement estimates may decrease during the last feeding phase of heavy pigs. Mahan et al. (1998b) observed that body Ca:P ratio greatly increased from 75 to 145 kg. This is because body Ca is mainly present in bone tissue, whereas P is present in soft and hard tissues; in heavy pigs, Ca and P deposition largely occurs in skeletal tissue with a declining deposition of P in muscle.

The dietary energy concentration may vary for heavy finishing pigs because of their increased capacity to adjust feed intake to meet the energy requirement (Suarez-Belloch et al. 2013). More importantly, increased gut capacity allows heavy pigs to digest and utilize energy from fibrous feedstuffs more efficiently through hindgut fermentation (Just et al., 1983; Noblet and Shi, 1994; Zanfi et al. 2012). This provides swine producers with an opportunity to lower feed cost by feeding fibrous feed ingredients. Galassi et al. (2007) compared growth performance of pigs fed 0, 12 and 24% wheat bran diets (11.8, 14.4, and 17.2% NDF, respectively) over different BW ranges; ADG and feed efficiency were worsened from 44 to 70 kg, numerically impaired from 70 to 98 kg, but were unaffected from 98 to 176 kg when wheat bran was included in the diets. In another study where pigs were fed 0, 15, and 30% sugar beet pulp in diets (14.2,

15.8, and 20.9% NDF, respectively), Galassi et al. (2005) observed that increasing dietary fiber worsened ADG and feed efficiency of pigs from 106 to 120 kg BW, but had no effect on pigs from 120 to 170 kg BW. This observation was supported by the fact that pigs fed the 3 different diets had similar energy digestibility measured at 154 kg. However, pigs fed in the two studies above were restrictively fed at approximately 2.25 kg DM/d. Future studies are needed to examine the effects of dietary fiber on growth performance of heavy pigs with *ad libitum* feeding. In addition, it is important to realize that pigs fed in a university environment may respond differently to the increased dietary fiber compared with pigs raised in a commercial environment because the feed intake of commercial pigs is subject to other restrictive factors, such as stocking density and hygiene (De la Llata et al., 2001). Meanwhile, the negative impact of dietary fiber on carcass yield should also be considered as the magnitude of this effect may be enlarged in heavy pigs due to their increased gut volume.

Feed additives and feeding strategies have been developed to help mitigate the increased fat deposition in heavy finishing pigs. Feeding ractopamine HCl before marketing allows pigs to produce heavier and leaner carcasses with better gain rate and efficiency than untreated pigs (Apple et al., 2007). The efficacy of ractopamine HCl has been confirmed in pigs raised up to 136 kg (Carr et al., 2009; Peterson et al., 2015). Porcine somatotropin is also effective in promoting pig growth performance and carcass leanness (Johnston et al., 1993), and such effects appear to be more prominent in heavy pigs (Kanis et al., 1990). However, somatotropin is not approved to be used in swine in the U.S.

Limiting fat deposition in heavy pigs may also be achieved via feed restriction. Slightly decreased feed intake increases nutrient digestibility, improves the efficiency of energy utilization, and decreases the amount of dietary energy partitioned to fat deposition. Nieto et al. (2012) suggested that pigs allowed to consume 70 and 95% of *ad libitum* feed intake were able

to retain similar amounts of body protein when raised to 150 kg. This finding indicates that heavy pigs may not require *ad libitum* feeding to attain the maximum protein deposition. Once pigs reach their genetic potential for maximum protein deposition, feed restriction becomes more effective in decreasing excessive fat gain. Although restricted feeding leads to decreased backfat thickness and slightly improved or unchanged G:F in heavy pigs, reduced ADG is often observed as a consequence of decreased feed intake (Hansson, 1974; García-Valverde et al., 2008; Kim et al. (2012). Moreover, feasibility of restricted feeding is questionable, at least in current U.S. production systems, with regards to the current feeder design and additional labor cost. As an alternative, feeding low-energy diets has been proposed to achieve the goal of restricting energy intake. However, the usefulness of this strategy is challenged by the fact that heavy finishing pigs increase feed intake to compensate for the reduced dietary energy density (Kim et al., 2005). It appeared that early finishing pigs fed low-energy diets had limited ability to adjust feed intake to maintain the same energy intake compared with pigs fed high-energy diets (Smith et al., 1999; Apple et al., 2004; Zhang et al., 2011), whereas heavy finishing pigs were able to maintain high feed and energy intake regardless of energy density of the diets (Suarez-Belloch et al., 2013). Although feeding low-energy diets effectively reduced backfat thickness, impaired ADG was still commonly observed. More importantly, inconsistent responses of caloric efficiency were often obtained when pigs were fed diets with decreased energy densities (Apple et al., 2004; Zhang et al., 2011; Suarez-Belloch et al., 2013), indicating a limited advantage of feeding low-energy diets to heavy finishing pigs.

Another challenge of raising heavy pigs is derived from the interactive effects between increasing marketing weight and gender on pig growth performance (Carr et al., 1978; Sather et al., 1980; Conte et al., 2011). Generally, barrows grow faster than gilts during late finishing phase, because gilts reach puberty at approximately 110 kg BW when declining feed intake and

growth rate are commonly observed (Hansson, 1974; Sather et al., 1980). Additionally, barrows have faster reductions of lean gain rate than gilts as body weights increase, indicating a different nutritional requirement for barrows and gilts. For instance, the Lys requirement suggested by the NRC (2012) growth model is approximately 0.05% lower for barrows than for gilts at both 130 and 140 kg BW. As a result, different feeding and marketing strategies are potentially needed for barrows and gilts. Through an economic model, Jolly et al. (1980) however argued that marketing both genders at equal weights resulted in negligible income penalty. Immunocastration has been used as an alternative of physical castration to eliminate boar taint while maintaining a pig growth performance similar to intact males. The efficacy of immunocastration has been verified for pigs with heavy marketing weight up to 176 kg (Zamaratskaia et al., 2008). However, as the length of mixed-housing period increases with marketing weight, it is possible that immunocastrated boars may stimulate the onset of puberty in gilts, but no research has been identified to address this question.

Animal housing

One major challenge of housing heavy pigs is the reduced floor space per pig. With a constant stocking density, space allowance becomes a limiting factor for ADFI and, subsequently, ADG of heavy pigs (Edmonds et al., 2003; Brumm, 2004; DeDecker et al., 2005). Weatherup et al. (1998) compared the growth performance of pigs housed individually and in groups (6 pigs/pen) and suggested that, with greater space allowance, individually housed pigs had a greater magnitude of increase in ADFI and less degree of reduction in ADG than group-housed pigs when marketing weight was raised. An allometric expression of the floor space required by pigs over a range of weights was proposed by Petherick (1983) and Baxter (1984) using the equation: $A, m^2 = k \times (BW, kg)^{0.667}$, where A represents floor space allowance and k represents a space allowance coefficient. When k is below 0.0336, decreased ADFI and ADG are often observed in

pigs housed on fully slatted floors (Gonyou et al., 2006). Calculations using the above equation with $k = 0.0336$, suggests that an average increment of $0.02 \text{ m}^2/\text{pig}$ is required for every 5 kg increase of pig BW from 125 to 150 kg in order not to decrease growth performance (Table 4). When adequate floor space cannot be provided, the impact of restricted pen space on pig performance is dependent on the magnitude of the restriction. A meta-analysis conducted by Flohr et al. (2015) established a set of equations to predict ADG, ADFI, and G:F based on pig BW. From this meta-analysis, for every 0.001 below the critical k value (0.0336), ADG, ADFI, and G:F are expected to decrease by 0.88, 0.58, and 0.31%, respectively, for pigs over 125 kg BW.

A pig removal strategy seems to be a good alternative to provide adequate floor space for heavy pigs in which the heaviest pigs within a pen are harvested first when they reach the target marketing weight, then the remainder pigs in the pen are provided increased floor space for improved growth. DeDecker et al. (2005) removed 25 and 50% of the heaviest pigs (13 or 26 out of 52 pigs/pen) when average pen weight reached 113 kg, which resulted in increased ADG (20.6 and 21.0%), ADFI (10.8 and 7.9%), and G:F (7.7 and 14.3%). Similarly, Jacela et al. (2009) observed that when 8 or 16% of the heaviest pigs (2 or 4 pigs out of a pen of 25) were removed when average pen weight reached 109 kg, pigs remaining in the pen had increased ADG (11.5 and 14.2%), ADFI (7.5 and 4.0%), and G:F (5.2 and 11.5%).

Appropriate feeder space is also essential for heavy pigs to maximize feed intake and gain. Excessive feeder space may increase feed wastage and worsen G:F when ample floor space is provided (Myers et al., 2012), whereas limiting feeder space harms growth performance especially when pigs have restricted floor space (Jungst et al., 2013). Size of a feeder hole should be 1.1 times the shoulder width (Brumm, 2012; Table 4), which can be estimated using: shoulder width (mm) = $64.0 \times (\text{BW, kg})^{0.33}$ (Petherick, 1983).

Height of waterers also should be adjustable based on the increased height of heavy pigs and the design of waterers. A general guideline for adjusting waterer height has been provided by Gonyou (1996). Nipple waterers pointed straight out from the wall should be placed at shoulder height, which can be predicted using: nipple waterer height, cm = $15 \times (\text{BW, kg})^{0.33}$. Nipple waterers mounted at a downwards angle should be placed 5 cm above the back of the pig, which can be estimated using: nipple waterer height, cm = $18 \times (\text{BW, kg})^{0.33}$. Finally, when water bowls are used, pigs should drink water with their head slightly lowered. Capacity of water pipes leading into the barn should also be sized accordingly to accommodate the increased total water consumption of heavier pigs. Nevertheless, excessive supply of water should be avoided in order to minimize water wastage and manure production. In addition, the height of pen partitions should be considered to accommodate the greater height of heavy pigs.

As BW increases, pigs generate more body heat but have decreased ability to dissipate this heat; thus, heavy pigs need lower critical ambient temperature and are more vulnerable to heat stress than light pigs (Renaudeau et al., 2011). According to a prediction equation from Brown-Brandt et al. (2004), heat production of pigs increases by 2% for every 5 kg increase in BW, indicating that barn ventilation rates need to be adjusted accordingly (Table 4). A production manual published by PIC (2014) recommends that barn temperature should be maintained at 16 °C for pigs from 96 to 138 kg and the minimal air exchange rates for pigs with 127 and 138 kg BW are 13.0 and 14.3 CFM/pig, respectively. In addition, ammonia emission is augmented as feed intake and manure production increase in heavy pigs (Ni et al., 2000), which can create a further challenge for proper barn ventilation.

Animal health

The duration of immunity following vaccinations for common swine pathogens when pigs are kept in barns to heavier weights is a complex subject. In theory, the need for vaccine

protection is decreased in heavier pigs because of their more developed immune system compared with young and naïve pigs. The necessity of providing heavy pigs an additional vaccination should be evaluated based on the immune status of the herd, because pigs with originally low antibody titers have greater response to vaccination, while pigs with originally high antibody titers have marginal benefits from the additional vaccination. It is also important to realize that the duration of immunity given by vaccination varies among vaccine products, types of vaccine (live vs. killed virus), and pathogens that vaccines are developed to against. Typically, vaccines designed to be given as 2 separate doses have longer protection than those given as single dose (Dick Hesse, personal communication). However, given the high economic cost of mortality in heavy pigs and the fact that risks of late-finishing disease, such as porcine reproductive and respiratory syndrome, influenza, and mycoplasma pneumonia, are still high, an additional dose of vaccine for heavy pigs has been occasionally used by producers (Dick Hesse, personal communication). However, for many vaccines, the effectiveness of an additional booster has not been critically evaluated and caution needs to be taken in regard to the legal withdraw period required following the vaccination.

Transportation

Transportation can induce a high amount of stress in heavy weight market pigs. Dalla Costa et al. (2009) observed elevated salivary cortisol concentrations and heart rate during loading and transport and Fitzgerald et al. (2009) reported higher mortality rate when pigs were transported at heavier weights compared with those marketed at lighter weights. As pigs grow heavier, they need more space provided in the trailer and better ventilation as they can become exhausted faster during transportation than light weight pigs. Meanwhile, the number of animals that can be transported per truck decreases with greater marketing weight (Table 4). Based on recommendations by Grandin (2012), truck space required by pigs transported during cool

weather increases from 0.43 to 0.50 m²/pig as marketing weight increases from 125 to 150 kg. Requirements for truck space may further increase when distance of transport and ambient temperature increase because pigs tend to spend more time laying (Guise et al., 1998; Torrey et al. 2013). The efficiency of loading and transporting heavy pigs also depends on the trailer design. Heavier pigs are reluctant to walk up a steep ramp and should be provided no more than a 15° ramp slope (Grandin, 2012).

Packing plant

With an increase in marketing weight, there are several practical packing plant considerations needed, including: processing equipment, transportation, and worker safety concerns. Through personal communication with meat scientists associated with large packing plants, increased body size, carcass length, and limb length of heavy pigs have been a main area of consideration. First, with an increase in final BW, line speed may decrease due to fewer numbers of pigs that can be stunned through carbon dioxide chambers used at nearly all major pork packing plants. Line speed can also be limited by USDA inspection, because a greater amount of time is needed to inspect a larger carcass. Second, as carcass length increases, pigs may not be able to be properly exsanguinated due to large variations in hind limb length and rail height. Rail height in older packing plants may also be a risk factor for de-hairing and scalding equipment as carcasses may drag on their backs at the bottom of scalding tanks. Furthermore, as pigs exit the de-hairing process, workers splitting carcasses will have to spin or roll a greater than 130 kg carcass into position. As the carcass continues through the harvesting process, longer limbs may also contribute to issues at the gambrel table, on conveyor belts, and on the main break table. Increased carcass weight may result in ergonomic concerns as workers need to handle and manipulate heavier hams, shoulders, and loins. Automated loin pullers and belly cutters may help mediate some of these issues. In addition, wind speeds and cooling times

required to properly chill heavy carcasses will need to be evaluated. Increased carcass size creates challenges on cooling capacity of packing plants, as greater airflow around and under the carcasses is needed. Coolers in older packing plants may already be running at the maximum wind speeds and cooling capacity and these packing plants may not have the capability to build additional cooling system. Finally, more storage space is also needed in coolers for the increased carcass weight and length.

Another consideration for increased carcass weights are consumer preferences. As carcass weight increases, there is a large weight increase in all of the primal cuts. Longer loins would be more desirable from a processing standpoint compared with increased loin diameter. This is because larger LM area would result in changes in portion controlled cutting. Chops cut to a standardized thickness would be heavier and resultantly more expensive during retail marketing, impacting the number of chops sold per package. Conversely, chops cut to a standardized weight would be thinner, requiring modifications to cooking methods currently used by both foodservice and consumers. It is unclear what impact these changes in chop thickness and weight would have on consumer preference. Furthermore, increasing marketing weight also affects the processing capacity of cull plants that specialize in handling lightweight cull pigs. When marketing weight range increases, cull pig weights would also have to increase. Some of these plants would have to drastically alter their plant design and space to process larger carcasses.

Conclusion

Many production variables are affected with increasing marketing weight. Generally, heavy weight market pigs eat more, but gain more slowly and less efficiently than pigs marketed at lighter weights. Heavier carcasses are associated with greater carcass yield, length, and LM area, but they also have greater backfat thickness and decreased percentage fat-free lean. Genetic

selection of lean-type pigs and research on nutritional requirements for pigs greater than 140 kg are needed to mitigate the reduction in feed efficiency and carcass leanness. Increasing marketing weight may result in minimal impacts on pork quality, but future studies are in need to evaluate consumer preferences on pork from heavy pigs with a focus on color, portion sizes, and sensory characteristics. In conclusion, as marketing weight increases approximately 0.5 kg per year (NASS, 2014), adjustments for nutritional and management guidelines, facility design, and packing plant equipment are necessary to accommodate increased biological and physical requirements of heavy weight market pigs.

LITERATURE CITED

- Apple, J., C. Maxwell, D. Brown, K. Friesen, R. Musser, Z. Johnson, and T. Armstrong. 2004. Effects of dietary lysine and energy density on performance and carcass characteristics of finishing pigs fed ractopamine. *J. Anim. Sci.* 82:3277-3287.
- Apple, J. K., P. J. Rincker, F. K. Mc, S. N. Keith, T. A. Carr, P. A. S. Armstrong, and P. D. Matzat. 2007. Review: Meta-analysis of the Ractopamine response in finishing swine. *Prof. Anim. Sci.* 23:179–196.
- Beattie, V., R. Weatherup, B. Moss, and N. Walker. 1999. The effect of increasing carcass weight of finishing boars and gilts on joint composition and meat quality. *Meat Sci.* 52:205-211.
- Baxter, S. 1984. Space and place. *Intensive pig production: environmental management and design.* Granada Publishing Ltd., London, UK. p. 210–254.
- Benz, J. M., M. D. Tokach, S. S. Dritz, J. L. Nelssen, J. M. DeRouchey, R. C. Sulabo, and R. D. Goodband. 2011. Effects of dietary iodine value product on growth performance and carcass fat quality of finishing pigs. *J. Anim. Sci.* 89:1419-1428.
- Bertol, T. M., E. A. Oliveira, A. Coldebella, V. L. Kawski, A. J. Scandolera, and M. B. Warpechowski. 2015. Meat quality and cut yield of pigs slaughtered over 100 kg live weight. *Arq. Bras. Med. Vet. Zootec.* 67:1166-1174.
- Brown-Brandt, T. M., J. A. Nienaber, H. Xin, R. S. Gates. 2004. A literature review of swine heat production. *Transactions of the ASAE.* 47:259-270.
- Brumm, M. 2004. The effect of space allocation on barrow and gilt performance. *J. Anim. Sci.* 82:2460-2466.

- Brumm, M. 2012. Impact of heavy market weights on facility and equipment needs. Proc. Allen D. Leman Swine Conference. St. Paul, MN. p. 165-168.
- Cannata, S., T. E. Engle, S. J. Moeller, H. N. Zerby, A. E. Radunz, M. D. Green, P. D. Bass, and K. E. Belk. 2010. Effect of visual marbling on sensory properties and quality traits of pork loin. *Meat Sci.* 85:428-434.
- Carr, S., D. Hamilton, K. Miller, A. Schroeder, D. Fernández-Dueñas, J. Killefer, M. Ellis, and F. McKeith. 2009. The effect of ractopamine hydrochloride (paylean[®]) on lean carcass yields and pork quality characteristics of heavy pigs fed normal and amino acid fortified diets. *Meat Sci.* 81:533-539.
- Carr, T., L. Walters, and J. Whiteman. 1978. Carcass composition changes in growing and finishing swine. *J. Anim. Sci.* 47:615-621.
- Cisneros, F., M. Ellis, F. McKeith, J. McCaw, and R. Fernando. 1996. Influence of slaughter weight on growth and carcass characteristics, commercial cutting and curing yields, and meat quality of barrows and gilts from two genotypes. *J. Anim. Sci.* 74:925-933.
- Conte, S., L. Boyle, N. O'Connell, P. Lynch, and P. Lawlor. 2011. Effect of target slaughter weight on production efficiency, carcass traits and behaviour of restrictively-fed gilts and intact male finisher pigs. *Livestock Science.* 136:169-174.
- Corino, C., M. Musella, and J. Mourot. 2008. Influence of extruded linseed on growth, carcass composition, and meat quality of slaughtered pigs at one hundred ten and one hundred sixty kilograms of liveweight. *J. Anim. Sci.* 86:1850-1860.

- Correa, J. A., C. Gariépy, M. Marcoux, and L. Faucitano. 2008. Effects of growth rate, sex, and slaughter weight on fat characteristics of pork bellies. *Meat Sci.* 80:550-554.
- Crome, P., F. McKeith, T. Carr, D. Jones, D. Mowrey, and J. Cannon. 1996. Effect of ractopamine on growth performance, carcass composition, and cutting yields of pigs slaughtered at 107 and 125 kilograms. *J. Anim. Sci.* 74:709-716.
- Crovetto, G. M., Galassi, G., Rapetti, L., Tamburini, A., 1999. Carbon, nitrogen and energy balance of heavy pigs fed diets with restricted protein and energy content. In: *Proceedings of the 13th Animal Science and Production Association National Congress, Piacenza, Italy.* p. 537–539.
- dalla Costa, O. A., J. V. Ludke, A. Coldebella, J. D. Kich, Costa, Mateus José Rodrigues Paranhos da, L. Faucitano, J. V. Peloso, and D. Dalla Roza. 2009. Effect of pre-slaughter management on physiological parameters of heavy-weight female pigs. *Ciência Rural.* 39:852-858.
- De la Llata, M., S. S. Dritz, M. D. Tokach, R. D. Goodband, J. L. Nelssen, and T. M. Loughin. 2001. Effects of dietary fat on growth performance and carcass characteristics of growing-finishing pigs reared in a commercial environment. *J. Anim. Sci.* 79:2643-2650.
- DeDecker, J., M. Ellis, B. Wolter, B. Corrigan, S. Curtis, E. Parr, and D. Webel. 2005. Effects of proportion of pigs removed from a group and subsequent floor space on growth performance of finishing pigs. *J. Anim. Sci.* 83:449-454.
- Durkin, I., M. Dadic, D. Brkic, B. Lukic, G. Kusec, M. Mikolin, and I. Jerkovic. 2012. Influence of gender and slaughter weight on meat quality traits of heavy pigs. *Acta agriculturae Slovenica.* 3:211-214.

- Edmonds, M., and D. Baker. 2003. Effect of dietary protein fluctuations and space allocation on performance and carcass quality of growing-finishing pigs. *J. Anim. Sci.* 81(11): 2783-2791.
- Fitzgerald, R., K. Stalder, J. Matthews, S. Kaster, and A. Johnson. 2009. Factors associated with fatigued, injured, and dead pig frequency during transport and lairage at a commercial abattoir. *J. Anim. Sci.* 87:1156-1166.
- Flohr, J. R. 2015. Effects of vitamin D supplementation and floor space on pig performance. Phd dissertation. Kansas State University, Manhattan, KS.
- Franci, O., C. Pugliese, R. Bozzi, A. Acciaioli, and G. Parisi. 2001. The use of multivariate analysis for evaluating relationships among fat depots in heavy pigs of different genotypes. *Meat Sci.* 58:259-266.
- Galassi, G., S. Colombini, L. Malagutti, G. Crovetto, and L. Rapetti. 2010. Effects of high fibre and low protein diets on performance, digestibility, nitrogen excretion and ammonia emission in the heavy pig. *Anim. Feed Sci. Technol.* 161:140-148.
- Galassi, G., G. Crovetto, and L. Malagutti. 2005. Effect of beet pulp on growing performance, digestibility, N balance, and ammonia emission in the heavy pig. *Ital. J. Anim. Sci.* 4(Suppl 2):458-460.
- Galassi, G., L. Malagutti, and G. M. Crovetto. 2007. Performance, nitrogen balance and ammonia emission from slurry in pigs fed high fibre diets. *Ital. J. Anim. Sci.* 6:227-239.

- García-Valverde, R., R. Barea, L. Lara, R. Nieto, and J. Aguilera. 2008. The effects of feeding level upon protein and fat deposition in iberian heavy pigs. *Livestock Science*. 114:263-273.
- Gonyou, H. 1996. Water use and drinker management. *Prairie Swine Centre 1996 Ann. Rep.* Saskatoon, Saskatchewan, Canada. p.74-80.
- Gonyou, H., M. Brumm, E. Bush, J. Deen, S. Edwards, T. Fangman, J. McGlone, M. Meunier-Salaun, R. Morrison, and H. Spolder. 2006. Application of broken-line analysis to assess floor space requirements of nursery and grower-finisher pigs expressed on an allometric basis. *J. Anim. Sci.* 84:229-235.
- Gu, Y., A. Schinckel, J. Forrest, C. Kuei, and L. Watkins. 1991. Effects of ractopamine, genotype, and growth phase on finishing performance and carcass value in swine: I. growth performance and carcass merit. *J. Anim. Sci.* 69:2685-2693.
- Gu, Y., A. Schinckel, and T. Martin. 1992. Growth, development, and carcass composition in five genotypes of swine. *J. Anim. Sci.* 70:1719-1729.
- Guise, H. J., H. L. Riches, B. J. Hunter, T. A. Jones, P. D. Warriss, and P. J. Kettlewell. 1998. The effect of stocking density in transit on the carcass quality and welfare of slaughter pigs. *Meat Sci.* 50:439-446.
- Grandin, T. 2012. Welfare of pigs during transport. *Pork information gateway*. Available at: <http://porkgateway.org/resource/welfare-of-pigs-during-transport>. (Accessed March, 2016).
- Hahn, J. D., and D. H. Baker. 1995. Optimum ratio to lysine of threonine, tryptophan, and sulfur amino acids for finishing swine. *J. Anim. Sci.* 73:482-489.

- Hansson, I. 1974. Effect of sex and weight on growth, feed efficiency and carcass characteristics of pigs. 1. growth rate and feed efficiency of boars, barrows and gilts. Swed. J. Agric. Res. 4:209-218.
- Hansson, I., K. Lundström, and B. Malmfors. 1975. Effect of sex and weight on growth, feed efficiency and carcass characteristics of pigs, 2: Carcass characteristics of boars, barrows and gilts, slaughtered at four different weights. Swed. J. Agric. Res. 5:69:80.
- Huff-Lonergan, E., T. J. Baas, M. Malek, J. C. M. Dekkers, K. Prusa, and M. F. Rothschild. 2002. Correlations among selected pork quality traits. J. Anim. Sci. 80:617-627.
- Jacela, J., M. D. Tokach, J. M. DeRouchey, R. D. Goodband, J. L. Nelssen, and S. S. Dritz. 2009. Economic impact of removing pigs before marketing on the remaining pigs' growth performance. Agricultural Experiment Station and Cooperative Extension Service. Kansas State University, Manhattan. p. 262-269.
- Johnston, M., J. Nelssen, R. Goodband, D. Kropf, R. Hines, and B. Schrick. 1993. The effects of porcine somatotropin and dietary lysine on growth performance and carcass characteristics of finishing swine fed to 105 or 127 kilograms. J. Anim. Sci. 71:2986-2995.
- Jolly, R., A. Sather, R. Patterson, B. Sonntag, A. Martin, and H. Freeden. 1980. Alternative market weights for swine: production economics. J. Anim. Sci. 51:804-810.
- Jungst, S., N. Matthews, C. Booher, B. Fields, T. Dreadin, S. Tabor, J. Anderson, J. Martin, A. Williams, M. Jobin, A. Sosnicki, and E. Wilson. 2012a. Growth curves for PIC327L sired pigs fed diets with differing energy levels. In: PIC Tech Memo. Hendersonville, TN. ES50-344.

- Jungst, S., N. Matthews, C. Booher, B. Fields, T. Dreadin, S. Tabor, J. Anderson, J. Martin, A. Williams, M. Jobin, A. Sosnicki, and E. Wilson. 2012b. Growth curves for commercial PIC337RG pigs fed high and low energy diets. In: PIC Tech Memo. Hendersonville, TN. ES51-344.
- Jungst, S., B. Fields, N. Matthews, M. Semler, C. Booher, and T. Dreadin. 2013. Effects of stocking density and feeder space allowance on growth and carcass quality of commercial pigs sired by PIC359 boars. In: *PIC Tech Memo*. Hendersonville, TN. ES067.
- Just, A., J. A. Fernandez, and H. Jørgensen. 1983. The net energy value of diets for growth in pigs in relation to the fermentative processes in the digestive tract and the site of absorption of the nutrients. *Livest. Prod. Sci.* 10:171–186.
- Kanis, E., G. Nieuwhof, K. De Greef, W. Van der Hel, M. Verstegen, J. Huisman, and P. Van der Wal. 1990. Effect of recombinant porcine somatotropin on growth and carcass quality in growing pigs: Interactions with genotype, gender and slaughter weight. *J. Anim. Sci.* 68:1193-1200.
- Kim, Y., S. Kim, M. Weaver, and C. Lee. 2005. Increasing the pig market weight: World trends, expected consequences and practical considerations. *Asian-Aust. J. Anim. Sci.* 18: 590-600.
- Latorre, M., E. García-Belenguer, and L. Ariño. 2008. The effects of sex and slaughter weight on growth performance and carcass traits of pigs intended for dry-cured ham from teruel (spain). *J. Anim. Sci.* 86:1933-1942.
- Latorre, M., R. Lázaro, D. Valencia, P. Medel, and G. Mateos. 2004. The effects of gender and slaughter weight on the growth performance, carcass traits, and meat quality characteristics of heavy pigs. *J. Anim. Sci.* 82:526-533.

- Latorre, M., P. Medel, A. Fuentetaja, R. Lázaro, and G. Mateos. 2003. Effect of gender, terminal sire line and age at slaughter on performance, carcass characteristics and meat quality of heavy pigs. *Anim. Sci.* 77:33-46.
- Leach, L., M. Ellis, D. Sutton, F. McKeith, and E. Wilson. 1996. The growth performance, carcass characteristics, and meat quality of halothane carrier and negative pigs. *J. Anim. Sci.* 74:934-943.
- Lo Fiego, D. P., P. Santoro, P. Macchioni, and E. De Leonibus. 2005. Influence of genetic type, live weight at slaughter and carcass fatness on fatty acid composition of subcutaneous adipose tissue of raw ham in the heavy pig. *Meat Sci.* 69:107-114.
- Mahan, D., and R. Shields. 1998a. Essential and nonessential amino acid composition of pigs from birth to 145 kilograms of body weight, and comparison to other studies. *J. Anim. Sci.* 76:513-521.
- Mahan, D., and R. Shields. 1998b. Macro-and micromineral composition of pigs from birth to 145 kilograms of body weight. *J. Anim. Sci.* 76:506-512.
- Mancini, R. A., and M. C. Hunt. 2005. Current research in meat color. *Meat Sci.* 71:100-121.
- Manini, R., A. Piva, A. Prandini, A. Mordenti, G. Piva, and J. Dourmad. 1997. Protein retention in italian heavy pigs: development of a factorial approach for the determination of lysine requirement. *Livest. Prod. Sci.* 47:253-259.
- Martin, A., A. Sather, H. Fredeen, and R. W. Jolly. 1980. Alternative market weights for swine. II. carcass composition and meat quality. *J. Anim. Sci.* 50:699-705.

- Moon, S. S., A. M. Mullen, D. J. Troy, H. S. Yang, S. T. Joo, and G. B. Park. (2003). Effect of pig slaughter weight on pork quality. *Korean J. Food Sci. Ani. Resour.* 23:315-320.
- Myers, A., R. Goodband, M. Tokach, S. Dritz, J. DeRouchey, and J. Nelssen. 2012. The effects of feeder adjustment and trough space on growth performance of finishing pigs. *J. Anim. Sci.* 90:4576-4582.
- NASS. 2014. *Agricultural Statistics*. USDA. National Agricultural Statistics Service. US Government Printing Office, Washington, D.C.
- Neely, J., R. Johnson, and L. Walters. 1979. Efficiency of gains and carcass characteristics of swine of two degrees of fatness slaughtered at three weights. *J. Anim. Sci.* 48:1049-1056.
- Ni, J., A. J. Heber, T. T. Lim, C. A. Diehl, R. K. Duggirala, B. L. Haymore, and A. L. Sutton. 2000. Ammonia emission from a large mechanically-ventilated swine building during warm weather. *J. Environ. Qual.* 29:751-758.
- Nieto, R., L. Lara, R. Barea, R. García-Valverde, M. Aguinaga, J. Conde-Aguilera, and J. Aguilera. 2012. Response analysis of the iberian pig growing from birth to 150 kg body weight to changes in protein and energy supply. *J. Anim. Sci.* 90:3809-3820.
- Noblet, J., and X. S. Shi. 1994. Effect of body weight on digestive utilization of energy and nutrients of ingredients and diets in pigs. *Livest. Prod. Sci.* 37:323–338.
- NRC. 2012. *Nutrient Requirements of Swine*. 11th rev. ed. Natl. Acad. Press, Washington, DC.
- Park, B., and C. Lee. 2011. Feasibility of increasing the slaughter weight of finishing pigs. *J. Anim. Sci. Technol.* 53:211-222.

Petherick, J. C. 1983. A biological basis for the design of space in livestock housing. In: Baxter, SH, MR Baxter and JASC Mac- Cormack (eds.) Farm Animal Housing and Welfare. Martinus Nijhoff Publisher, Boston. p. 103-120.

Peloso, J., P. Lopes, L. Gomide, S. Guimarães, and P. Carneiro. 2010. Carcass and ham quality characteristics of heavy pigs from different genetic groups intended for the production of dry-cured hams. *Meat Sci.* 86:371-376.

Peterson, C., C. Pilcher, H. Rothe, J. Marchant-Forde, M. Ritter, S. Carr, C. Puls, and M. Ellis. 2015. Effect of feeding ractopamine hydrochloride on growth performance and responses to handling and transport in heavy-weight pigs. *J. Anim. Sci.* 93:1239-1249.

Piao, J., J. Tian, B. Kim, Y. Choi, Y. Kim, and I. K. Han. 2004. Effects of sex and market weight on performance, carcass characteristics and pork quality of market hogs. *Asian-Aust. J. Anim. Sci.* 17:1452-1458.

PIC. 2014. Wean to finish manual. Available at:

http://www.pic.com/Images/Users/1/salesportal/literature/manuals/04252014_WeanToFinish_Manual.pdf. (Accessed March, 2016)

Raj, St., G. Skiba, D. Weremko, F. Fandrejewski, W. Migdal, F. Borowiec, and E. Polawska. 2010. The relationship between the chemical composition of the carcass and the fatty acid composition of intramuscular fat and backfat of several pig breeds slaughtered at different weights. *Meat Sci.* 86:324-330.

Renaudeau, D., J. Gourdine, and N. St-Pierre. 2011. A meta-analysis of the effects of high ambient temperature on growth performance of growing-finishing pigs. *J. Anim. Sci.* 89:2220-2230.

- Sather, A., A. Martin, R. Jolly, and H. Fredeen. 1980. Alternative market weights for swine. I. feedlot performance. *J. Anim. Sci.* 51:28-36.
- Schinckel, A., S. Pence, M. Einstein, R. Hinson, P. Preckel, J. Radcliffe, and B. Richert. 2006. Evaluation of different mixed model nonlinear functions on pigs fed low-nutrient excretion diets. *The Professional Animal Scientist.* 22:401-412.
- Serrano, M., D. Valencia, A. Fuentetaja, R. Lázaro, and G. Mateos. 2008. Effect of gender and castration of females and slaughter weight on performance and carcass and meat quality of iberian pigs reared under intensive management systems. *Meat Sci.* 80:1122-1128.
- Shields, R., D. Mahan, and P. Graham. 1983. Changes in swine body composition from birth to 145 kg. *J. Anim. Sci.* 57:43-54.
- Shull, C. 2013. Modeling growth of pigs reared to heavy weights. Phd dissertation. University of Illinois at Urbana-Champaign, Urbana, IL.
- Smith, J., M. Tokach, P. O'Quinn, J. Nelssen, and R. Goodband. 1999. Effects of dietary energy density and lysine: Calorie ratio on growth performance and carcass characteristics of growing-finishing pigs. *J. Anim. Sci.* 77:3007-3015.
- Suarez-Belloch, J., M. Sanz, M. Joy, and M. Latorre. 2013. Impact of increasing dietary energy level during the finishing period on growth performance, pork quality and fatty acid profile in heavy pigs. *Meat Sci.* 93:796-801.
- Torrey, S., R. Bergeron, L. Faucitano, T. Widowski, N. Lewis, T. Crowe, J. Correa, J. Brown, S. Hayne, and H. Gonyou. 2013. Transportation of market-weight pigs: II. effect of season

- and location within truck on behavior with an eight-hour transport. *J. Anim. Sci.* 91:2872-2878.
- Virgili, R., M. Degni, C. Schivazappa, V. Faeti, E. Poletti, G. Marchetto, M. Pacchioli, and A. Mordenti. 2003. Effect of age at slaughter on carcass traits and meat quality of italian heavy pigs. *J. Anim. Sci.* 81:2448-2456.
- Wagner, J., A. Schinckel, W. Chen, J. Forrest, and B. Coe. 1999. Analysis of body composition changes of swine during growth and development. *J. Anim. Sci.* 77:1442-1466.
- Weatherup, R., V. Beattie, B. Moss, D. Kilpatrick, and N. Walker. 1998. The effect of increasing slaughter weight on the production performance and meat quality of finishing pigs. *Anim. Sci.* 67:591-600.
- Zamaratskaia, G., L. Rydhmer, H. K. Andersson, G. Chen, S. Lowagie, K. Andersson, and K. Lundström. 2008. Long-term effect of vaccination against gonadotropin-releasing hormone, using ImprovacTM, on hormonal profile and behaviour of male pigs. *Anim. Reprod. Sci.* 108:37-48.
- Zanfi, C., and M. Spanghero. 2012. Digestibility of diets containing whole ear corn silage for heavy pigs. *Livestock Science.* 145:287-291.
- Zhang, G., X. Yi, L. Chu, L. Ning, J. Htoo, and S. Qiao. 2011. Effects of dietary net energy density and standardized ileal digestible lysine: Net energy ratio on the performance and carcass characteristic of growing-finishing pigs fed low crude protein supplemented with crystalline amino acids diets. *Agricultural Sciences in China.* 10:602-610.

Table 1. Summary of studies investigating the effects of market weight on overall growth performance (changes per 10 kg marketing weight increase)¹

Reference	Initial weight, kg	Marketing weight, kg	Pigs/pen	Space/pig, m ²	Total pigs	ADG, g	ADFI, g	G:F
Neely et al. (1979)	15	100,113,127	6	-	200	8.7	52.7	-0.004
Sather et al. (1980)	2	73,84,98,109,123,134	4	1.44	288	-16.0	102.0	-0.015
Kanis et al. (1990)	60	100,140	1	-	96	-19.5	56.3	-0.012
Johnston et al. (1993)	59	105,127	3	2.30	120	8.0	54.0	-0.003
Cisneros et al. (1996)	60	100,115,130,145,160	4	1.17	160	4.0	100.0	-0.006
Leach et al. (1996)	40	110,125,140	4	1.20	144	-18.6	-	-0.010
Weatherup et al. (1998) ²	50	92,103,113,125	1	6.00	96	-9.2	111.3	-0.017
Weatherup et al. (1998) ³	50	92,103,113,125	6	1.00	288	2.8	91.9	-0.014
Latorre et al. (2003)	25	122,136	5	1.10	240	7.1	78.6	-0.009
Latorre et al. (2004)	75	116,124,133	8	1.00	192	-38.0	-3.0	-0.010
Piao et al. (2004)	27	100,110,120,130	4	1.01	224	-7.3	76.4	-0.014
Latorre et al. (2008)	107	120,125,130,135,140	10	1.05	200	-54.9	-78.0	-0.010
Serrano et al. (2008)	25	145,156	15	1.50	360	8.2	163.6	-0.013
Shull (2013) Exp.2	6	113,125,136,147,159,170,181	20	1.06	2240	-3.6	58.1	-0.012
Average ⁴	-	-	-	-	-	-4.0	78.1	-0.011

¹ Generated by simple linear regression analyses by EXCEL.

² Individual housing was evaluated.

³ Group housing was evaluated.

⁴ Studies by Latorre et al. (2004 and 2008) were excluded from the calculation because pigs were reported to be under heat stress; study by Serrano et al. (2008) was excluded from calculation due to the use of Iberian obese pig breed that was uncommonly used in north America pig production.

Table 2. Summary of studies investigating the effects of marketing weight on carcass characteristics (changes per 10 kg marketing weight increase)¹

Reference	Marketing weight, kg	Yield, %	Backfat, mm	Fat-free lean, %	LM area, cm ²	Length, cm	Subprimal yield, %			
							Belly	Loin	Shoulder	Ham
Hansson (1975)	68,88,108,128	0.84	2.1	-1.03	1.7	3.1	-	-	-	-
Carr et al. (1978)	45,68,91,114,136	-	2.0	-1.00	2.2	2.4	-	-	-	-0.09
Neely et al. (1979)	100,113,127	-	1.0	0.07	2.0	1.9	-	-	-	-
Sather et al. (1980) and Martin et al. (1980)	73,84,98,109,123,134	-	-	-0.47	2.3	2.3	0.53	-	-0.48	-0.20
Shields et al. (1983)	56,76,90,107,127,146	1.05	2.8	-	1.7	2.3	0.12	-0.19	-0.15	-0.28
Kanis et al. (1990)	100,140	-	1.1	-0.55	-	-	-	-	-	-
Gu et al. (1991 and 1992)	100,114,127	0.34	3.0	-1.09	1.1	2.3	-	-	-	-
Johnston et al. (1993)	105,127	0.05	0.9	-0.18	2.7	-	-	-	-	-
Crome et al. (1996)	107,125	0.33	2.1	-	1.2	2.1	0.61	-0.18	-	0.14
Cisneros et al. (1996)	100,115,130,145,160	0.32	1.6	-	1.8	1.9	0.09	0.40	-0.18	-0.16
Leach et al. (1996)	110,125,140	0.16	1.4	-1.59	-0.02	1.7	0.45	-0.38	0.08	-0.19
Weatherup et al. (1998) ²	92,103,113,125	0.68	1.6	-1.28	-	-	-	-	-	-
Weatherup et al. (1998) ³	92,103,113,125	0.35	1.5	0.09	-	-	-	-	-	-
Beattie et al. (1999)	96,108,121,133	0.29	-	-	2.2	-	-	-	-	-
Wagner et al. (1999)	25,45,64,84,100,129,152	0.67	2.3	-0.77	2.3	2.7	-	-0.09	-	-0.19
Latorre et al. (2003)	122,136	0.29	0.5	-	-	2.1	-	-0.21	-0.21	-0.36
Virgili et al. (2003)	144,182	0.34	-	-	1.5	-	-	-0.29	-0.32	-0.19
Latorre et al. (2004)	116,124,133	0.77	2.9	-	-	2.4	-	-	-0.29	0.04
Piao et al. (2004)	100,110,120,130	-0.49	0.9	0.05	2.3	3.1	-	-	-	-
Correa et al. (2008)	107,115,125	0.41	-	-	-	2.0	0.13	-0.12	0.12	-0.28
Corino et al. (2008)	111,160	0.38	2.0	-1.85	-	-	-	-0.06	-	-
Latorre et al. (2008)	120,125,130,135,140	0.48	2.5	-	-	1.3	-	-0.18	-0.02	-0.34
Serrano et al. (2008)	145,156	0.91	1.2	-	-	-	-	-0.18	0.09	0.36
Shull (2013) Exp.1	75,91,106,121,134,147,168	-	1.7	-	2.6	-	-	-	-	-
Shull (2013) Exp.2	115,124,134,145,157,166,176	0.43	1.8	-1.36	1.9	-	-	-	-	-
Average ⁴	-	0.41	1.8	-0.78	1.8	2.2	0.32	-0.13	-0.16	-0.17

¹ Generated by simple linear regression analyses by EXCEL.

² Individual housing was evaluated.

³ Group housing was evaluated.

⁴ Study by Serrano et al. (2008) was excluded from calculation due to the use of Iberian obese pig breed which was uncommonly used in north America pig production.

Table 3. Summary of studies investigating the effects of marketing weight on pork quality (changes per 10 kg marketing weight increase)¹

Reference	Marketing weight, kg	L*	a*	b*	Initial pH	Ultimate pH	Drip loss, %	WBSF ² , kg
Beattie et al. (1999)	92, 105, 118, 131	0.52	-0.02	0.18	-	-0.01	0.22	-0.05
Bertol et al. (2015) ³	100, 115, 130, 145	-0.23	0.23	-	-0.05	0.01	0.34	-
Bertol et al. (2015) ⁴	100, 115, 130, 146	0.04	0.16	-	-0.04	-	0.08	0.14
Cisneros et al. (1996)	100, 115, 130, 145, 160	-	-	-	-0.01	-0.02	0.29	-0.08
Durkin et al. (2012)	120, 130, 140, 150, 160, 170	-0.14	0.34	0.10	-0.02	-0.01	0.27	0.01
Leach et al. (1996)	110,125,140	-1.23	0.30	-0.14	-0.01	-	-0.35	0.24
Latorre et al. (2004)	116, 124, 133	-2.48	-	-0.24	-	-	-	0.11
Moon et al. (2003)	95, 105, 115, 125	-	-	-	-	-0.04	0.21	-
Piao et al. (2004)	100, 110, 120, 130	1.15	1.18	0.42	-	0.02	-4.75	-0.04
Virgili et al. (2003) ⁵	144,182	-0.01	0.10	-0.17	-0.01	-0.05	-	0.16
Virgili et al. (2003) ⁶	144,182	-	-	-	-	-	-0.34	-
Weatherup et al. (1998)	92,103,113,125	0.17	0.12	0.20	-	-0.01	0.30	-
Average		-0.25	0.30	0.05	-0.02	-0.01	-0.11	0.06

¹ Generated by simple linear regression analyses by EXCEL.

² Warner-Bratzler Shear Force.

³ Ham was evaluated.

⁴ Longissimus dorsi was evaluated.

⁵ Semimembranosus was evaluated.

⁶ Resulted due to 20.7% drip loss in 100 kg pigs; no differences in methodology present.

⁷ Study by Piao et al. (2004) was excluded from calculation for drip loss effect due to the abnormally high value reported (greater than 3 standard deviations from the mean of all values).

Table 4. Changes in facility recommendations for pigs based on final marketing weight

Items	Marketing weight, kg					
	125	130	135	140	145	150
Floor space/pig ¹ , m ²	0.84	0.86	0.89	0.91	0.93	0.95
Feeder space ² , cm	34.6	35.1	35.5	36.0	36.4	36.8
Drinker height, cm						
Right-angled waterer ³	73.8	74.8	75.7	76.6	77.5	78.4
Downward waterer ⁴	88.6	89.7	90.8	91.9	93.0	94.1
Heat production ⁵ , kcal/h	242.1	248.1	254.0	259.7	265.5	271.1
Pigs/truck ⁶	163	156	151	145	140	136
Truck space/pig ⁷ , m ²	0.43	0.44	0.45	0.47	0.48	0.50

¹ Estimated using: floor space, m² = $k \times (\text{BW, kg})^{0.667}$, where $k = 0.0336$ (Gonyou et al., 2006).

² Estimated using: feeder space = $1.1 \times \text{shoulder width}$ (Brumm, 2012), and shoulder width, mm = $64.0 \times (\text{BW, kg})^{0.33}$ (Petherick, 1983).

³ Estimated using: right-angled waterer height, cm = $15 \times (\text{BW, kg})^{0.33}$ (Gonyou, 1996).

⁴ Estimated using: downward waterer height, cm = $18 \times (\text{BW, kg})^{0.33}$ (Gonyou, 1996).

⁵ Estimated using: heat production (W/kg) = $14.11 \times (\text{BW, kg})^{-0.38}$ (Brown-Brandt et al., 2004)

⁶ Assuming maximum truck load of 20321.1 kg.

⁷ Adapted from recommendation from Grandin (2012).