

**Title:** Sources of and adequacy of protein intake among pregnant women in the United States – **NPB #19-046**

**Investigator:** Mary M. Murphy

**Institution:** Exponent, Inc.

**Date Submitted:** February 1, 2021

**Industry Summary:** Limited information is available on protein intake and adequacy of intake among pregnant women in the United States (U.S.). Protein intake by women of childbearing age in the U.S. is generally assumed to be adequate, though requirements for protein increase during the second and third trimesters of pregnancy and limited data suggest that some women may fail to meet recommended intakes during this critical time. The purpose of this study was to assess protein intake and adequacy of intake among pregnant women in the U.S. Using data from a nationally representative sample of 528 pregnant women, mean usual intake of protein was estimated by trimester of pregnancy. Findings from this study indicate that approximately one in eight women in the second and third trimesters of pregnancy have protein intakes below the recommended levels. Results from the study also show that percent protein intake from animal sources was positively associated with likelihood of meeting trimester-specific protein recommendations. Guidance for pregnant women should include advice to consume adequate levels of protein from a variety of protein sources, including animal proteins for women following a Healthy U.S.-Style Dietary Pattern. Consumption of animal proteins, including lean meats, can help women meet protein requirements during pregnancy.

E-mail: [mmurphy@exponent.com](mailto:mmurphy@exponent.com); Telephone: 202-772-4953

### **Key Findings:**

- Approximately one in eight women in the second and third trimesters of pregnancy in the U.S. have protein intakes below the recommended levels.
- Percent protein intake from animal sources was positively associated with likelihood of women meeting trimester-specific protein recommendations.
- Guidance for pregnant women should include advice to consume adequate levels of protein from a variety of protein sources, including animal proteins for women following a Healthy U.S.-Style Dietary Pattern.

**Keywords:** dietary intake; nutrient intake adequacy; macronutrient; protein; maternal diet; pregnancy

---

These research results were submitted in fulfillment of checkoff-funded research projects. This report is published directly as submitted by the project's principal investigator. This report has not been peer-reviewed.

---

**For more information contact:**

**National Pork Board** • PO Box 9114 • Des Moines, IA 50306 USA • 800-456-7675 • Fax: 515-223-2646 • [pork.org](http://pork.org)

---

**Scientific Abstract:** Limited information is available on protein intake and adequacy of protein intake among pregnant women. Using data from a sample of 528 pregnant women in the National Health and Nutrition Examination Surveys (NHANES) 2003-2012, usual intake of protein (g/day and g/kg body weight (bw)/day) and prevalence of intake below the Estimated Average Requirement (EAR) by trimester of pregnancy were calculated using the National Cancer Institute method. Percent contributions to protein intake by source (i.e., plant and animal, including type of animal source) were also calculated. Mean usual intake of protein was  $88 \pm 4.3$ ,  $82 \pm 3.1$ , and  $82 \pm 2.9$  g/day among women in trimester 1, 2, and 3 of pregnancy, respectively, or  $1.30 \pm 0.10$ ,  $1.35 \pm 0.06$ , and  $1.35 \pm 0.05$  g/kg bw/day, respectively. An estimated 4.5% of women in the first trimester of pregnancy consumed less protein than the EAR of 0.66 g/kg bw/day; among women in the second and third trimesters of pregnancy, 12.1% and 12.8% of women, respectively, consumed less protein than the EAR of 0.88 g/kg bw/day. Animal sources of protein accounted for approximately 65% of total protein. Findings from this study show that one in eight women in the second and third trimesters of pregnancy have inadequate intake of protein. Pregnant women should be encouraged to consume sufficient levels of protein from a variety of sources.

**Introduction:** The importance of nutrition during pregnancy for both maternal and child health outcomes is well recognized [1]. Nutrient needs increase during pregnancy to support fetal growth and development along with alterations in maternal metabolism and tissue development [2]. Requirements for micronutrients increase during periconception and during pregnancy, while requirements for energy and protein increase particularly during the second and third trimesters [2,3].

Protein requirements during pregnancy were derived by the Institute of Medicine (IOM) using a factorial method that accounts for protein deposition and the additional of protein required to support healthy weight gain during pregnancy [3]. In the first trimester of pregnancy, the additional protein needs are minimal at approximately 1 g/day, and the estimated average requirement (EAR) is not different from the value of 0.66 g/kg bw/day for non-pregnant females [3]. In the second and third trimesters of pregnancy, protein needs increase an average of 21 g/day, resulting in an EAR of 0.88 g/kg bw/day [3]. Assuming a standard body weight for women, these EARs are equivalent to a daily protein intake of 38 g in the first trimester and 59 g in the latter two trimesters. The recommended dietary allowance (RDA) for protein during the first trimester of pregnancy is estimated at 46 g/day (0.8 g/kg bw/day), and at 71 g/day (1.1 g/kg bw/day) during the second and third trimesters [3].

Protein requirements established outside North America using the factorial method are similar to recommendations established by the IOM. The European Food Safety Authority (EFSA) identified additional protein needs of 1, 9, and 28 g/day, respectively by trimester of pregnancy [4], while D-A-CH, the nutrition societies of Germany, Austria, and Switzerland, set nutrient requirements for protein at 0.8, 0.9, and 1.0 g/kg bw/day during the first, second and third trimesters, respectively [5]. The indicator amino acid oxidation (IAAO) method has been studied as an alternative method to estimate protein requirements, including requirements during pregnancy, and tends to identify higher protein requirements relative to the balance or factorial methods (+15-73%) [5-7].

Nationally representative nutrient intake estimates for pregnant women in the U.S. are limited, and the available assessments of intake adequacy have largely focused on micronutrient intakes [8]. Using the National Health and Nutrition Examination Survey (NHANES) 2001-2014, mean usual intake of protein by pregnant, non-lactating women in the U.S. was estimated at 81.9 g/day, though prevalence of protein intake below the EAR was not examined [8]. Based on data from 2007-2010, mean usual intake of protein by pregnant women in the U.S. was estimated at 78.6 g/day [9]; distributions of usual intake of protein among the

sample indicated that the EAR for women in the second and third trimester of pregnancy fell between the 5th and 10th percentiles of intake, suggesting greater inadequacy of protein intake among women during pregnancy relative to inadequacy among non-pregnant women [10,11].

The purpose of this study was to assess protein intake and adequacy of intake among a representative sample of pregnant women in the U.S. population. To provide further insight on protein intake, sources of protein (i.e., plant and animal, including type of animal source) consumed by pregnant women were characterized.

**Objectives:** The overall objective of this study is to assess protein intake and adequacy of protein intake among pregnant women in the United States (US) using data from a nationally representative sample of pregnant women, and to examine a possible association between protein intake adequacy and dietary protein intake from the consumption of animal sources. Secondary aims of the study are to (a) characterize the contributions of protein by (i.e., plant and animal, including red meat overall and specifically beef, pork, as well as poultry, eggs, and seafood) to total protein intake; (b) characterize the contributions of protein by source of protein (i.e., plant and animal) consumed by pregnant women by status of protein adequacy; and (c) characterize gestational weight gain pattern by status of protein adequacy and source of protein (i.e., plant and animal).

### **Materials & Methods:**

Data Source and Study Population: This cross-sectional study was conducted with data collected in NHANES and the dietary recall component known as What We Eat in America (WWEIA) [12,13]. The NHANES provides nationally representative nutrition and health data used to develop prevalence estimates for nutrition and health status measures in the U.S. and serves as the foundation of many nutrient assessments for national nutrition policy [12]. Each NHANES cycle includes an in-person household interview, a health examination in a mobile examination center (MEC), and a telephone follow-up interview. Approval for the NHANES data collection was provided by the National Center for Health Statistics (NCHS) Research Ethics Review Board.

Pregnancy status was assessed with a urine pregnancy test during the MEC examination and women self-identified as pregnant reported month of pregnancy. More recent NHANES cycles have not included the question on month of pregnancy, thus this analysis was limited to survey cycles 2003-2004, 2005-2006, 2007-2008, 2009-2010, and 2011-2012. The sample population for this study was pregnant women ages 20-44 years as identified by a positive urine pregnancy test, self-identified month of pregnancy, were not breastfeeding, and provided two valid dietary recalls as determined by NCHS. The final sample of pregnant, non-lactating women included in this analysis is 528.

Population Characteristics: Demographic, lifestyle, and reproductive health characteristics were included to characterize the sample population by trimester of pregnancy. Usual intake of energy by trimester of pregnancy was also estimated. Self-reported month of pregnancy was categorized as trimester 1 (months 1, 2, 3), trimester 2 (months 4, 5, 6), and trimester 3 (months  $\geq 7$ ). Demographic characteristics include age at screening, race/ethnicity (non-Hispanic white, non-Hispanic black, Mexican American or other Hispanic, or other race), poverty income ratio (PIR;  $<1.85$ ,  $\geq 1.85$ ), education status (less than high school, high school diploma, some college, undergraduate degree or higher), and marital status (married, widowed or divorced or separated, never married). Lifestyle characteristics include self-reported smoking status (never smoked, past smoker, current smoker), physical activity ( $<10$  minutes/week, 10 to 150 minutes/week,  $\geq 150$  minutes/week), and self-reported dietary supplement use (Yes, No). Reproductive health characteristics include parity, defined as the

number of live births, and gestational weight gain (GWG) classified as inadequate, adequate, or excessive using criteria established by the IOM [13]. Pre-pregnancy BMI was calculated from self-reported pre-pregnancy body weight collected during the in-home interview and standing height measured in the MEC.

Dietary Intake Data: The WWEIA component of NHANES consists of two 24-hour dietary recalls (midnight to midnight); the first dietary recall is collected during the MEC interview and the second is collected during the follow-up telephone interview 3-10 days after the MEC interview. Each recall is collected by trained interviewers using USDA's Automated Multiple-Pass method. For each food reported in WWEIA, the survey specific USDA Food and Nutrient Database for Dietary Studies (FNDDS) databases provide the translation of foods as reported consumed into gram amounts and ingredients for the processing of energy and nutrient intakes [14].

Assessment of Protein Intake and Adequacy: Estimates of protein intake were generated as grams per day (g/day) and as grams per kilogram pre-pregnancy body weight per day (g/kg bw/day) assuming actual body weight when it was within a normal healthy weight range (body mass index (BMI)  $\geq 18.5$  and  $< 24.9$  kg/m<sup>2</sup>), or the pre-pregnancy body weight that would place the woman at the nearest endpoint of the healthy range. Adjustment of body weight to a value aligning with a healthy body weight is consistent with the approach used in assessments of protein intake and adequacy of intake [15,16]. Protein intake adequacy was defined as the prevalence of usual intake of protein below the EAR.

Identification of Protein by Source: The USDA food codes representing each food consumed were disaggregated into the survey-specific Standard Reference (SR) items used by USDA to process nutrient values [17]. Each SR code was then categorized as plant or animal, with animal sources further categorized as red meat (i.e., beef, pork, other), poultry, cured meat & poultry, seafood, eggs, and dairy. Contributions of protein by source were calculated per 100 g food code. For SR codes representing mixtures of protein sources, contributions from different sources were estimated from USDA data by translating each SR code into food pattern equivalents [18]. Protein intake by source was summed across all foods consumed over the two days of recall per respondent and divided by the 2-day average protein intake per person to develop percent contributions per person.

Assessment of Protein Intake using Alternate Recommendations: As an exploratory analysis, adequacy of protein intake among pregnant women in the U.S. was also assessed using alternate reference values for adequate protein intake during pregnancy developed from the IAAO method. These alternate EAR correspond to intake of 1.22 g/kg bw/day during the early stages (~16 weeks) of pregnancy and 1.52 g/kg bw/day during late stages (~36 weeks) [6,7]. In this analysis, the alternate EAR during the first trimester of pregnancy was assumed to be 0.93 g/kg bw/day, which is the IAAO value for non-pregnant adults. The alternate EAR during the second trimester of pregnancy was assumed to be 1.22 g/kg bw/day, and the alternate EAR during the third trimester was assumed to be 1.52 g/kg bw/day. Consistent with the assessments with the IOM EARs, the pre-pregnancy body weight for each pregnant woman in the sample was adjusted to a body weight that would place the woman at the nearest endpoint of the healthy range ( $\geq 18.5$  to  $< 24.9$  kg/m<sup>2</sup>).

Statistical Analyses: Usual intakes of protein and energy were estimated using the National Cancer Institute (NCI) method [19] and the SAS macros developed by NCI for modeling a single dietary component with covariates for day of the week (weekday/weekend) and sequence of the dietary recall (day 1 or 2). Balanced repeated replicate weights (Fay adjustment factor = 0.3) based on 2-day dietary recall statistical weights were used for SE estimation [20].

Prevalence of usual intake of protein below the IOM EAR and the alternate IAAO EAR reference values for protein adequacy [6,7] were calculated using the cut-point method [21]. Using the 2-day average intakes, mean percent contributions of protein intake from plant and animal sources were estimated for all pregnant women and by subcategory of trimester of pregnancy. Logistic regression models were used to test the hypothesis that percent of total protein consumed as animal protein is associated with an increased odds of meeting the trimester-specific RDA for protein and increased likelihood of adequate GWG. In the analysis of GWG, women were classified in two categories: adequate and inadequate (including excess) GWG. The logistic regression models were initially adjusted for the following covariates: age, trimester of gestation, parity, race/ethnicity, marital status, education, family poverty income ratio, energy intake, pre-pregnancy body mass index, smoking status, and physical activity. Non-statistically significant covariates were sequentially dropped from the models. Protein intake was represented by the 2-day average intake in the regression analyses. Statistical analyses were completed using SAS (version 9.4, SAS Institute Inc., Cary, NC, USA) and STATA V12.1 (StataCorp, College Station, Texas 77845, USA). All analyses used the appropriate statistical weights provided in NHANES to account for oversampling, survey non-response, and post-stratification and estimates of the standard errors (SEs) and confidence intervals were design adjusted. Estimates are presented as mean  $\pm$  SE. Statistical significance was defined as Bonferroni-adjusted  $p < 0.05$ .

## **Results:**

Protein Characteristics: Of the 528 pregnant women included in this sample, 109 women were in trimester 1 of pregnancy, 207 were in trimester 2, and 212 were in trimester 3. With the exception of parity and GWG, population characteristics did not vary by trimester of pregnancy. The population was primarily non-Hispanic white, married women, a PIR  $\geq 1.85$ , at least some college education, and 1 or more previous deliveries (Table 1). The majority of pregnant women achieved at least 10 minutes of moderate physical activity per day, were non-smokers, and reported intake of dietary supplements in the past 30 days.

Protein Intake and Adequacy of Protein Intake: Usual intake of protein was  $88 \pm 4.3$ ,  $82 \pm 3.1$ , and  $82 \pm 2.9$  g/day among women in trimester 1, 2, and 3 of pregnancy, respectively (Table 2), or  $1.30 \pm 0.10$ ,  $1.35 \pm 0.06$ , and  $1.35 \pm 0.05$  g/kg bw/day based on pre-pregnancy body weight in a healthy range.

Adequacy of protein intake was assessed by calculating the percentage of women with usual intake of protein below the EAR. An estimated  $4.5 \pm 4.8\%$  of women in the first trimester of pregnancy consumed less than the EAR of 0.66 g/kg bw/day; given the sample of 109 women in the first trimester of pregnancy, this estimate is potentially less statistically reliable [22,23]. Among women in the second and third trimesters of pregnancy,  $12.1 \pm 4.3\%$  and  $12.8 \pm 4.6\%$  of women, respectively, consumed less than the EAR of 0.88 g/kg bw/day.

Intake of Protein by Source: Approximately two thirds ( $65.2 \pm 1.0\%$ ) of total protein intake was from animal sources and ranged from  $67.0 \pm 1.7\%$ , to  $63.9 \pm 1.8\%$ , and  $65.0 \pm 1.0\%$  for women in the first, second, and third pregnancy trimester, respectively. Red meat, poultry, and cured meat & poultry combined accounted for approximately one-third of total intake of protein among pregnant women ( $34\%$ ) while dairy accounted for  $22.8 \pm 0.9\%$  of protein intake. Protein from eggs and seafood each accounted for approximately 4% of protein consumed by pregnant women. No consistent trend across the three pregnancy trimesters was observed in the contribution of the various sources to total protein intake (Table 3).

Protein by Source and Protein Recommendations: After dropping the non-statistically significant covariates, the final logistic regression model evaluated the association between adequacy of protein intake relative to the RDA and percent protein from animal sources, and included

covariates for pregnancy trimester, race/ethnicity, total energy intake, and pre-pregnancy BMI category. The odds ratio associated with percent protein from animal sources in the final model was  $1.1 \pm 0.02$  (p-value  $<0.05$ ), indicating that pregnant women were more likely to meet the trimester-specific protein RDA as their percent protein intake from animal sources increased. Women with higher total energy intake and those in the first pregnancy trimester were also more likely to meet the protein RDA than those with lower energy intake or those in pregnancy trimester two or three.

Protein by Source and Gestational Weight Gain: Logistic regression models were also used to evaluate the association between the likelihood of adequate GWG and meeting the protein RDA and percent protein from animal sources. Women were classified in two categories: adequate and inadequate (including excess) GWG, and the models included the following covariates: age, trimester of gestation, parity, race/ethnicity, marital status, education, family poverty income ratio, energy intake, pre-pregnancy body mass index, smoking status, and physical activity. Since the analyses summarized above indicated a statistically significant association between meeting the protein RDA requirement and percent protein from animal sources, two separate analyses of adequate GWG were conducted using each of these predictors separately. Non-statistically significant covariates were sequentially dropped from the model. No statistically significant associations between adequate GWG and either meeting the protein RDA or percent protein from animal sources were found in the final models. However, in both models, women classified as having a normal pre-pregnancy BMI were more likely have adequate GWG, and those with higher total energy intake were less likely to have adequate GWG.

Assessment of Protein Adequacy using Alternate Requirements: Adequacy of protein intake was also assessed based on alternate protein requirements derived for pregnant women using the IAAO method [6,7]. Protein intake among pregnant women was below the alternate EAR for  $18.5 \pm 10.0\%$ ,  $40.1 \pm 5.5\%$ , and  $66.9 \pm 5.1\%$  of pregnant women in trimester 1, 2, and 3, respectively (Figure 1).

**Discussion:** In this study, protein intake and adequacy of protein intake were examined in a nationally representative sample of pregnant women by trimester of pregnancy. The prevalence of inadequate protein intake among women in the first trimester of pregnancy was approximately 5% based on the EAR of 0.66 g/kg bw/day. Among women in the second or third trimester of pregnancy, 12-13% of women consumed inadequate levels of protein based on the EAR of 0.88 g/kg bw/day. Results from this study indicate that many pregnant women, in particular those beyond the first trimester of pregnancy, consumed suboptimal levels of protein relative to recommendations established by the IOM.

The estimates of protein intake and adequacy of intake developed in this study are based on data collected in 2003-2012. While more recent NHANES data are available, data collected in recent NHANES cycles do not include information on trimester of pregnancy, consequently it would not be possible to compare intakes to trimester-specific requirements. A recent review of protein intake from USDA shows that total protein intake among U.S. females ages 20-39 years remained relatively unchanged between 2005-2006 (74 g/day) and 2015-2016 (73 g/day), and that the estimated contribution of animal sources to total protein intake was approximately 67% [24], which is consistent with the estimated contributions calculated in the current study. The stable intake of protein over the last decade and the consistent contribution of protein from animal sources suggest that the estimates of intake developed in this study may provide relevance for current protein intake and adequacy among pregnant women. The data also provide a point of reference for comparison in future analyses.

The 2020-2025 Dietary Guidelines for Americans (DGA) are the first DGA to include nutrition recommendations for women during the life stages of pregnancy and lactation [25]. The DGAs

did not assess adequacy of protein intake in units of g/kg bw/day, though the DGA did examine intake of protein foods relative to dietary recommendations. Based on intakes in 2013-2016, the Dietary Guidelines Advisory Committee (DGAC) reported that nearly half (47%) of pregnant women consumed fewer ounce equivalents of protein foods than recommended, with particularly low intakes of seafood and plant-based protein [26, 27]. However, the DGAC did not identify protein as a nutrient of concern for pregnant women. Results in the current study show that percent protein intake from animal sources was positively associated with likelihood of meeting the trimester-specific RDA.

The assessment of protein adequacy based on alternate requirements developed with the IAAO method indicate that approximately 40-67% of pregnant women consumed inadequate levels of protein during periods of significant fetal and maternal protein deposition. Adequacy of protein intake based on requirements derived with the IAAO method therefore suggest that suboptimal protein intake among pregnant women may be of greater concern than indicated by assessment with the IOM EAR. Further research on protein needs during pregnancy is needed.

Strengths of this study include a relatively large, nationally representative sample of pregnant women with identification by trimester of pregnancy that permitted comparisons of protein intake to trimester-specific requirements. Protein intake and adequacy of intake were based on usual intake estimates derived using established statistical methods. Percent contributions of protein by source were based on 2-day average intakes, which may provide a more representative profile of typical intake than a single day of recall.

As previously noted, the data used in this analysis are not current, though evidence among non-pregnant women suggests that protein intake has not changed over the last decade. Other limitations of the analysis must be acknowledged. Estimates of protein intake on a body weight basis were calculated using self-reported pre-pregnancy weights, as measured pre-pregnancy body weight values are not available in NHANES due to the cross-sectional nature of the survey. Pregnant women missing the trimester of pregnancy were excluded from the sample population which could introduce bias. Additionally, dietary recalls were self-reported and therefore subject to misreporting.

Findings from this study indicate that approximately one in eight women in the second and third trimesters of pregnancy have protein intakes below the EAR. Guidance for pregnant women should include advice to consume adequate levels of protein from a variety of protein sources. For women following a Healthy U.S.-Style Dietary Pattern, protein foods to encourage include lean meats, poultry, eggs, seafood, and plant-based proteins.

**Table 1.** Characteristics of sample of pregnant women ages 20-44 years by trimester of pregnancy in 2003-2012 National Health and Nutrition Examination Survey

Characteristic <sup>1</sup>	Total (n=528)	Trimester 1 (n=109)	Trimester 2 (n=207)	Trimester 3 (n=212)
Age (years)	28.7 ± 0.44	27.9 ± 0.61	28.3 ± 0.69	29.6 ± 0.74
Race/ethnicity (%)				
Mexican American/ Other Hispanic	23.1 ± 3.26	24.2 ± 7.06	22.2 ± 4.76	23.2 ± 3.91
Non-Hispanic white	52.9 ± 4.25	58.6 ± 8.86	47.3 ± 5.87	53.7 ± 5.84
Non-Hispanic black	13.5 ± 2.68	8.7 ± 3.69	20.9 ± 4.41	10.6 ± 3.15
Other race	10.5 ± 2.52	8.6 ± 4.13	9.7 ± 3.92	12.5 ± 4.15
Poverty income ratio (%)				
<1.85	34.6 ± 3.92	32.6 ± 8.73	38.3 ± 5.49	32.8 ± 5.36
≥1.85	65.4 ± 3.92	67.4 ± 8.73	61.7 ± 5.49	67.2 ± 5.36
Education status (%)				
< High school	19.9 ± 2.47	14.1 ± 3.94	23.4 ± 5.28	20.8 ± 4.7
High school diploma	13.7 ± 1.93	9.7 ± 3.19	15.1 ± 3.68	15.4 ± 3.28
Some college	33.2 ± 3.34	47.7 ± 8.15	29.7 ± 4.69	26.3 ± 5.25
Undergraduate degree or higher	33.2 ± 3.36	28.5 ± 7.18	31.9 ± 5.55	37.5 ± 6.39
Parity (%)				
0	25.6 ± 3.55	31.7 ± 7.94	30.0 ± 5.53	18.3 ± 2.97
1	43.5 ± 3.82	40.4 ± 7.1	51.8 ± 6.67	38.8 ± 6.1
≥2	30.9 ± 3.2	27.9 ± 5.78	18.2 ± 3.24	42.9 ± 5.62
Marital status (%)				
Married	69.3 ± 3.06	68.6 ± 6.74	68.9 ± 5.29	70.2 ± 5.00
Widowed/divorced/separated	2.3 ± 0.69	3.7 ± 1.84	1.0 ± 0.43	2.3 ± 1.25
Never married	28.4 ± 3.1	27.7 ± 6.62	30.1 ± 5.12	27.5 ± 4.98
Physical activity, minutes/week (%)				
< 10	31.7 ± 3.44	24.9 ± 6.41	34.4 ± 5.82	34.1 ± 5.86
10 to <150	30.1 ± 3.57	35.2 ± 6.83	29.1 ± 4.97	27.5 ± 5.38
≥ 150	38.2 ± 3.94	39.9 ± 7.26	36.5 ± 5.76	38.4 ± 6.74
Smoking status (%)				
Never smoked	69.3 ± 3.53	69.5 ± 6.73	65.5 ± 6.33	72.4 ± 5.16
Past smoker	23.3 ± 3.46	21.3 ± 6.15	26.7 ± 5.94	21.7 ± 4.72
Current smoker	7.4 ± 1.48	9.2 ± 3.04	7.7 ± 3.25	6.0 ± 2.67
Dietary supplement use (%)	85.6 ± 2.11	75.7 ± 5.76	88.6 ± 3.35	89.7 ± 3.79
Pre-pregnancy BMI (%)				
Underweight	5.2 ± 1.7	4.1 ± 3.45	5.8 ± 2.41	5.4 ± 3.27
Normal weight	46.7 ± 3.57	40.6 ± 8.21	39.3 ± 5.31	57.0 ± 5.68
Overweight	21.8 ± 2.91	16.3 ± 6.07	25.7 ± 5.67	22.0 ± 4.16
Obese	26.3 ± 3.35	39.0 ± 8.11	29.1 ± 4.86	15.7 ± 3.56
GWG (%)				
Inadequate	19.7 ± 2.87	35.8 ± 8.47	16.0 ± 3.78	12.2 ± 3.63
Adequate	31.5 ± 3.87	30.8 ± 8.45	30.4 ± 5.86	32.8 ± 6.62
Excess	48.9 ± 3.75	33.3 ± 7.47	53.6 ± 6.24	55.1 ± 6.73
Usual energy intake (kcal/day)	2272 ± 66.0	2275 ± 107.2	2298 ± 79.7	2247 ± 96.3

Values reported as mean ± SE.

<sup>1</sup>Estimates based on total sample in each trimester category, excluding missing data for PIR (n=27), parity (n=19), physical activity (n=2), pre-pregnancy BMI status (n=9), and GWG status (n=9). Statistically significant differences across trimester of pregnancy for parity and GWG category (p<0.05).



**Table 2.** Usual protein intake among pregnant women ages 20-44 years in 2003-2012 National Health and Nutrition Examination Survey

Trimester of Pregnancy	Percentile							
	Mean	5 <sup>th</sup>	10 <sup>th</sup>	25 <sup>th</sup>	50 <sup>th</sup>	75 <sup>th</sup>	90 <sup>th</sup>	95 <sup>th</sup>
grams protein per day (g/day)								
Total (n=528)	84 ± 2.3	57 ± 3.8	62 ± 3.3	71 ± 2.7	83 ± 2.3	95 ± 2.9	107 ± 4.2	115 ± 5.2
1 (n=109)	88 ± 4.3	61 ± 5.4 <sup>1</sup>	66 ± 5.2 <sup>1</sup>	75 ± 4.8	87 ± 4.4	100 ± 4.7	112 ± 5.5 <sup>1</sup>	120 ± 6.3 <sup>1</sup>
2 (n=207)	82 ± 3.1	55 ± 4.0	60 ± 3.6	70 ± 3.2	80 ± 3.0	92 ± 3.8	104 ± 5.1	111 ± 6.0
3 (n=212)	82 ± 2.9	56 ± 4.1	61 ± 3.7	70 ± 3.2	82 ± 2.9	94 ± 3.5	105 ± 4.9	112 ± 5.9
grams protein per kilogram body weight (g/kg bw/day) <sup>2</sup>								
Total (n=528)	1.34 ± 0.04	0.71 ± 0.10	0.82 ± 0.09	1.04 ± 0.07	1.31 ± 0.04	1.61 ± 0.05	1.88 ± 0.08	2.06 ± 0.10
1 (n=109)	1.30 ± 0.10	0.68 ± 0.15 <sup>1</sup>	0.79 ± 0.14 <sup>1</sup>	1.00 ± 0.13	1.27 ± 0.11	1.57 ± 0.09	1.84 ± 0.09 <sup>1</sup>	2.03 ± 0.09 <sup>1</sup>
2 (n=207)	1.35 ± 0.06	0.71 ± 0.09	0.84 ± 0.08	1.06 ± 0.06	1.32 ± 0.05	1.61 ± 0.08	1.90 ± 0.11	2.07 ± 0.13
3 (n=212)	1.35 ± 0.05	0.72 ± 0.09	0.83 ± 0.08	1.05 ± 0.06	1.33 ± 0.05	1.62 ± 0.07	1.89 ± 0.11	2.07 ± 0.13

Values reported as mean ± SE.

<sup>1</sup> Values may be less statistically reliable based on guidance from NCHS [22] and reported variance inflation factors up to 2.51 in NHANES 2005-2012 [23].

<sup>2</sup> Intake of protein as g/kg bw/day calculated from pre-pregnancy body weight assuming reported body weight when it was within a normal healthy weight range (BMI ≥18.5 and <24.9), or the weight that would place the woman at the nearest endpoint of the healthy range.

**Table 3.** Mean percent of protein intake by source among pregnant women ages 20-44 years in 2003-2012 National Health and Nutrition Examination Survey


Protein Source		Total (n=528)	Trimester 1 (n=109)	Trimester 2 (n=207)	Trimester 3 (n=212)
Animal	Total	65.2 ± 1.0	67.0 ± 1.7	63.9 ± 1.8	65.0 ± 1.0
	<i>Total red meat<sup>1</sup></i>	15.0 ± 1.1	17.5 ± 2.2	11.5 ± 1.3	16.2 ± 1.7
	Beef	12.3 ± 1.0	14.3 ± 2.3	10.0 ± 1.2	13.0 ± 1.6
	Pork	2.3 ± 0.5	2.0 ± 0.6	1.5 ± 0.4	3.2 ± 1.0
	Poultry	13.2 ± 1.1	12.5 ± 2.0	13.6 ± 1.5	13.4 ± 1.9
	Cured meat & poultry	5.8 ± 0.6	5.5 ± 0.9	6.2 ± 1.0	5.8 ± 1.1
	Dairy	22.8 ± 0.9	21.3 ± 1.7	24.9 ± 1.4	22.0 ± 1.3
	Egg	4.4 ± 0.4	5.5 ± 1.0	3.9 ± 0.5	4.0 ± 0.5
	Seafood	4.0 ± 0.7	4.7 ± 1.9	4.0 ± 1.1	3.7 ± 0.9
Plant	Total	34.8 ± 1.0	33.0 ± 1.7	36.1 ± 1.8	35.0 ± 1.0

Values reported as mean percent ± SE.

<sup>1</sup>Includes intake from other red meat for three women.

No statistically significant (p<0.05) differences between groups.

**Figure 1.** Percent of pregnant women ages 20-44 years with protein intake below reference requirements, 2003-2012 National Health and Nutrition Examination Survey



Values presented as means. Protein requirements - IOM EAR: 0.66 g/kg bw/day for trimester 1, 0.88 g/kg bw/day for trimesters 2 and 3 [3]; IAAO EAR: 0.93 g/kg bw/day for trimester 1, 1.22 g/kg bw/day for trimester 2, 1.52 g/kg bw/day for trimesters 3 [6,7]. Abbreviations: EAR – estimated average requirement, IAAO - indicator amino acid oxidation, IOM – Institute of Medicine.

## References

1. Procter, S.B.; Campbell, C.G. Position of the Academy of Nutrition and Dietetics: nutrition and lifestyle for a healthy pregnancy outcome. *J Acad Nutr Diet* **2014**, *114*, 1099-1103.
2. Picciano, M.F. Pregnancy and lactation: physiological adjustments, nutritional requirements and the role of dietary supplements. *J Nutr* **2003**, *133*, 1997S-2002S.
3. Institute of Medicine. Dietary Reference Intakes for Energy, Carbohydrates, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids. Washington, DC: National Academy Press: 2005.
4. EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies). Scientific Opinion on Dietary Reference Values for protein. *EFSA Journal* **2012**, *10*, 2557.
5. Richter M; Baerlocher K; Bauer JM; Elmadfa I; Heseker H; Leschik-Bonnet E; Stangl G; Vokert D; Stehle P. Revised Reference Values for the Intake of Protein. *Ann Nutr Metab*, **2019**, *74*, 242-250.
6. Elango, R.; Ball, R.O. Protein and Amino Acid Requirements during Pregnancy. *Adv Nutr* **2016**, *7*, 839S-844S.
7. Stephens, T.V.; Payne, M.; Ball, R.O.; Pencharz, P.B.; Elango, R. Protein requirements of healthy pregnant women during early and late gestation are higher than current recommendations. *J Nutr* **2015**, *145*, 73-78.
8. Bailey, R.L.; Pac, S.G.; Fulgoni, V.L., 3rd; Reidy, K.C.; Catalano, P.M. Estimation of Total Usual Dietary Intakes of Pregnant Women in the United States. *JAMA Netw Open* **2019**, *2*, e195967.
9. U.S. Department of Agriculture; Agricultural Research Service. Protein (g/day): Mean intake and percentiles of usual intake from food and beverages by Dietary Reference Intake age-gender groups in the United States, females 19-50, 2007-2010. 2014. Available online: <https://health.gov/our-work/food-nutrition/previous-dietary-guidelines/2015/advisory-report/appendix-e-2/appendix-e-24#protein> (accessed 01 March 2020).
10. Berryman, C.E.; Lieberman, H.R.; Fulgoni, V.L., 3rd; Pasiakos, S.M. Protein intake trends and conformity with the Dietary Reference Intakes in the United States: analysis of the National Health and Nutrition Examination Survey, 2001-2014. *Am J Clin Nutr* **2018**, *108*, 405-413.
11. Fulgoni, V.L., 3rd. Current protein intake in America: analysis of the National Health and Nutrition Examination Survey, 2003-2004. *Am J Clin Nutr* **2008**, *87*, 1554S-1557S.
12. Ahluwalia, N.; Dwyer, J.; Terry, A.; Moshfegh, A.; Johnson, C. Update on NHANES Dietary Data: Focus on Collection, Release, Analytical Considerations, and Uses to Inform Public Policy. *Adv Nutr* **2016**, *7*, 121-134.
13. Institute of Medicine. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC: National Academy Press: 2009.
14. U.S. Department of Agriculture; Agricultural Research Service. Food and Nutrient Database for Dietary Studies. Available online: <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fndds-download-databases/> (accessed 01 March 2020).
15. Moshfegh A; Goldman J; Cleveland L. What We Eat in America, NHANES 2001-2002: Usual Nutrient Intakes from Food Compared to Dietary Reference Intakes. U.S. Department of Agriculture, Agricultural Research Service. 2005. Available online: <https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/0102/usualintaketables2001-02.pdf> (accessed 01 March 2020).
16. U.S. Department of Agriculture, A.R.S. Usual Nutrient Intakes, WWEIA NHANES 2013-2016. 2019. Available online: [https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/usual/Usual\\_Intake\\_gender\\_WWEIA\\_2013\\_2016.pdf](https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/usual/Usual_Intake_gender_WWEIA_2013_2016.pdf) (accessed 01 March 2020).

17. U.S. Department of Agriculture; Agricultural Research Service. USDA National Nutrient Database for Standard Reference, Legacy Release. Available online: <https://data.nal.usda.gov/dataset/usda-national-nutrient-database-standard-reference-legacy-release> (accessed 01 March 2020).
18. U.S. Department of Agriculture; Agricultural Research Service. Food Patterns Equivalents Database. Available online: <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fped-databases/> (accessed 01 March 2020).
19. Tooze, J.A.; Kipnis, V.; Buckman, D.W.; Carroll, R.J.; Freedman, L.S.; Guenther, P.M.; Krebs-Smith, S.M.; Subar, A.F.; Dodd, K.W. A mixed-effects model approach for estimating the distribution of usual intake of nutrients: the NCI method. *Stat Med* **2010**, *29*, 2857-2868.
20. Herrick, K.A.; Rossen, L.M.; Parsons, R.; Dodd, K.W. Estimating Usual Dietary Intake From National Health and Nutrition Examination Survey Data Using the National Cancer Institute Method. *Vital Health Stat 2* **2018**, 1-63.
21. Institute of Medicine. Dietary Reference Intakes: Applications in Dietary Assessment. Washington, DC: National Academy Press: 2000.
22. National Center for Health Statistics. Analytic and Reporting Guidelines: The Third National Health and Nutrition Examination Survey, NHANES III (1988-94). Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 1996. Available online: <https://www.n.cdc.gov/nchs/data/nhanes/analyticguidelines/88-94-analytic-reporting-guidelines.pdf> (accessed 24 January 2021).
23. U.S. Department of Agriculture; Agricultural Research Service. WWEIA Data Tables. Available online: <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/wweia-data-tables/> (accessed 01 March 2020).
24. Hoy MK; Clemens JC; Moshfegh A. Protein Intake of Adults in the U.S.: What We Eat in America, NHANES 2015-2016. Food Surveys Research Group Dietary Data Brief No. 29. January 2021, revised slightly from July 2020. 2021. Available online: [https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/DBrief/29\\_Protein\\_Intake\\_of\\_Adults\\_1516.pdf](https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/DBrief/29_Protein_Intake_of_Adults_1516.pdf) (accessed on 26 January 2021).
25. U.S. Department of Agriculture; U.S. Department of Health and Human Services. Dietary Guidelines for Americans, 2020-2025.9th Edition. 2020. Available online: [https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary\\_Guidelines\\_for\\_Americans\\_2020-2025.pdf](https://www.dietaryguidelines.gov/sites/default/files/2020-12/Dietary_Guidelines_for_Americans_2020-2025.pdf) (accessed 24 January 2021).
26. Dietary Guidelines Advisory Committee. Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services. U.S. Department of Agriculture, Agricultural Research Service, Washington, DC: 2020.
27. 2020 Dietary Guidelines Advisory Committee and Data Analysis Team. Data Supplement for Food Group and Nutrient Distribution: All Life Stages . 2020 Dietary Guidelines Advisory Committee Project. Washington, DC: U.S. Department of Agriculture and U.S. Department of Health and Human Services., 2020. Available online: [https://www.dietaryguidelines.gov/sites/default/files/2020-07/DA\\_Supplement\\_FoodGroup\\_NutrientDistribution.pdf](https://www.dietaryguidelines.gov/sites/default/files/2020-07/DA_Supplement_FoodGroup_NutrientDistribution.pdf) (accessed 24 January 2021).